University of Vermont

UVM ScholarWorks

College of Arts and Sciences Faculty
Publications

College of Arts and Sciences

Summer 7-7-2007

Insertion of Benzylisocyanide into a Zr-P bond and Rearrangement. Atom-Economical Synthesis of a Phosphaalkene

Samantha N. MacMillan

Vassar College, samacmillan@gmail.com

Joseph M. Tanski

Vassar College, jotanski@vassar.edu

Rory Waterman

University of Vermont, rory.waterman@uvm.edu

Follow this and additional works at: <https://scholarworks.uvm.edu/casfac>

Part of the [Inorganic Chemistry Commons](#)

Recommended Citation

MacMillan SN, Tanski JM, Waterman R. Insertion of benzyl isocyanide into a Zr-P bond and rearrangement. Atom-economical synthesis of a phosphaalkene. *Chem Commun (Camb)*. 2007 Oct 28;(40):4172-4. doi: 10.1039/b709506f. <http://pubs.rsc.org/en/content/articlepdf/2007/cc/b709506f>

This Article is brought to you for free and open access by the College of Arts and Sciences at UVM ScholarWorks. It has been accepted for inclusion in College of Arts and Sciences Faculty Publications by an authorized administrator of UVM ScholarWorks. For more information, please contact scholarworks@uvm.edu.

Insertion of Benzyliisocyanide into a Zr–P bond and Rearrangement. Atom-Economical Synthesis of a Phosphaalkene[†]

Samantha N. MacMillan,^a Joseph M. Tanski^a and Rory Waterman^{*b}

Receipt/Acceptance Data [DO NOT ALTER/DELETE THIS TEXT]

Publication data [DO NOT ALTER/DELETE THIS TEXT]

DOI: 10.1039/b000000x [DO NOT ALTER/DELETE THIS TEXT]

Reaction of (N₃N)ZrPPh (1; N₃N = N(CH₂CH₂NSiMe₃)₃)³⁻ with PhCH₂N≡C afford the 1,1-insertion product (N₃N)Zr[C(PhPh)=NCH₂Ph] (2), which thermally rearranges to the phosphaalkene-containing complex, (N₃N)Zr[N(CH₂Ph)C(H)=PPh] (3).

An intensified interest in phosphaalkenes has emerged due to the advantageous properties the P=C bond displays as a synthetic precursor,¹ transition-metal ligand,² and part of conjugated materials.³ In typical syntheses of phosphaalkenes, transmetalation or elimination reactions are commonly employed,⁴ and steric protection is often required to kinetically trap the P=C bond.⁵ As part of our on-going interest in developing new metal-mediated reactions that form phosphorus–element bonds,⁶ we wish to report an unusual transformation that forms a new phosphalkene with perfect atom-economy.

Reaction of (N₃N)ZrPPh (1; N₃N = N(CH₂CH₂NSiMe₃)₃)³⁻ with 1 equiv of benzyliisocyanide in benzene solution at ambient temperature afforded the 1,1-insertion product, (N₃N)Zr[C(PhPh)=NCH₂Ph] (2, eqn 1).[‡] Complex 2 was isolated as a pale-orange powder by lyophilization of the benzene solution and characterized by NMR (¹H, ¹³C, and ³¹P) and infrared spectroscopy. The phosphine proton of 2 displays a significantly downfield shifted resonance at δ 6.038 and a large coupling to phosphorus, *J*_{PH} = 259 Hz, compared to other (N₃N)Zr-species containing a phenylphosphide moiety.⁶ This assignment was confirmed by observation of equivalent P–H coupling in the ³¹P NMR spectrum of 2. Additionally, ν_{PH} = 2280 cm⁻¹ was observed in the infrared. The strong P–H scalar coupling may arise from increased s-character of the P–H bond resulting from some delocalization of the phosphorus lone pair into the imine π-system, a feature that may also be responsible for the downfield chemical shift of the phosphorus proton. Other spectroscopic features support the formulation given, including an imine ν_{CN} = 1705 cm⁻¹ in the infrared and an imine carbon resonance at δ 263.2 with *J*_{PC} = 99.5 Hz in the ¹³C NMR spectrum of 2.

Insertion of unsaturated organic molecules into metal-phosphido bonds is well known since the first example, insertion of CO into the Hf–P bond of Cp*HfCl₂(P^tBu₂), by Bercaw and coworkers.⁷ More recently, Hey-Hawkins and coworkers have reported the

insertion of phenylisocyanide into the Zr–P bond of Cp*₂ZrClP(SiMe₃)₂ (Cp* = C₅MeH₄)⁸. Insertion reactions of other unsaturated substrates into Zr–P bonds have also appeared.⁹

Complex 2 exhibits limited thermal stability and gradually decomposes even as a solid when stored in the dark at –30 °C under N₂. The decomposition of 2 is significantly accelerated in solution and is qualitatively more rapid in polar solvents such as Et₂O rather than benzene or toluene. The product of the decomposition, in all cases, is the phosphaalkene-containing complex, (N₃N)Zr[N(CH₂Ph)C(H)=PPh] (3, eqn 2). Complex 3 was prepared directly by reaction of phosphido 2 with benzyliisocyanide in benzene followed by heating.[‡] Observation of the reaction by ¹H and ³¹P NMR showed quantitative conversion to 3, which was isolated in 78% yield as analytically pure, pale-yellow crystals from a concentrated Et₂O solution. A similar rearrangement of the phenylisocyanide insertion product, Cp*₂ZrCl[η²-N(Ph)=CP(SiMe₃)₂], that Hey-Hawkins reported (*vide supra*), was not observed, possibly due to the η²-coordination of the imine or the lower propensity of substituents other than hydride to engage in migrations.⁸

Spectroscopic properties of 3 support the formulation given. Notably, the phosphaalkene carbon resonates at δ 193.1 with *J*_{PC} = 55.6 Hz in the ¹³C NMR spectrum, and the phosphorus nucleus resonates at δ 91.9 with no primary P–H scalar coupling in the ³¹P NMR spectrum of 3—values consistent with a phosphaalkene moiety. The hydrogen atom of the phosphaalkene carbon resonates at δ 10.29 and is broad (Δν_{1/2} ~ 40 Hz) at ambient temperature in the ¹H NMR spectrum. Investigation of this complex by variable temperature ¹H and ³¹P NMR spectroscopy (215 – 320 K) in toluene-*d*₈ solution revealed a dynamic process with an activation barrier Δ*G*[‡] = 15.1 kcal/mol. At lower (<280 K) and higher (>320 K) temperatures, the P–H coupling becomes resolved (*J*_{PH} ~ 12

^a Department of Chemistry, Vassar College, Poughkeepsie, NY, 12604, USA.

^b Department of Chemistry, University of Vermont, Burlington, VT 05405, USA. Fax: 01 802 656-8705; Tel: 01 802 656-0278; E-mail: rory.waterman@uvm.edu

[†] Electronic Supplementary Information (ESI) available: Complete experimental, spectroscopic, and analytical details and X-ray data for 3 (CCDC number: XX). See <http://dx.doi.org/10.1039/b000000x/>

Hz), suggesting that the process is a thermal *E/Z* isomerization of the phosphalkene (eqn 3).¹⁰

The structure of **3** was confirmed by a single crystal X-ray diffraction study, and a perspective view of the complex is shown in Figure 1.⁸ The phosphalkene is in the *E* configuration in the solid state, N and C(16) are planar, and the phenyl substituent of phosphorus is coplanar with P and C(16), implicit of an sp^2 -hybridized phosphorus center and delocalization. The P–C bond length of 1.716(2) Å is somewhat long compared to other structurally characterized phosphalkenes.^{4,5} This feature may result from a zwitterionic resonance contributor where there is some C–N double bond character and a delocalized system. In support of this hypothesis, the C–N bond length of 1.362(2) Å is slightly shorter than expected for a C–N single bond. Known amine-substituted phosphalkenes display similar bond lengths.¹¹ Interestingly, complex **3** is a rare instance of metal complex containing a phosphalkene moiety that is *not* involved in coordination to the metal center.¹²

Figure 1. Perspective view of the molecular structure of **3** with hydrogen atoms except H(16) omitted for clarity. Thermal ellipsoids shown at the 35% probability level. Selected metrical parameters: N(5)–C(16) = 1.362(2), N(5)–C(17) = 1.465(2), P–C(16) = 1.716(2), P–C(21) = 1.831(2), C(16)–H(16) = 0.96(2), Zr–N(1) = 2.064(1), Zr–N(3) = 2.071(1), Zr–N(2) = 2.081(1), Zr–N(5) = 2.187(1), Zr–N(4) = 2.498(1) Å; C(16)–N(5)–C(17) = 114.3(1), C(16)–N(5)–Zr = 121.75(9), C(17)–N(5)–Zr = 122.12(9), N(5)–C(16)–P = 126.4(1), N(5)–C(16)–H(16) = 114.2(1), P–C(16)–H(16) = 119.3(1), C(16)–P–C(21) = 101.63(7)°.

A rich reaction chemistry has developed around phosphalkenes as facile syntheses evolved.^{1,4–5} There are several common routes to these molecules including 1,2-elimination, condensation, and rearrangement reactions.⁴ Synthesis of phosphalkenes by a 1,2-hydride migration appears not to have been previously reported. However, the most related reaction in the literature is a 1,2-phenyl migration from a P(V) intermediate.¹³

This synthesis, insertion of an organic isocyanide in to the Zr–P bond of a primary phosphido complex followed by rearrangement, takes advantage of phenylphosphine as the phosphorus source and commercially available benzyliisocyanide. Such a strategy avoids salt elimination and exhibits perfect atom economy. We are currently expanding this novel transformation into a general synthesis.

This work was supported by the University of Vermont and the U.S. National Science Foundation (Grant # 0521237 to JMT). The authors thank Prof. John Protasiewicz for helpful discussions.

Notes and references

- [†] *Experimental data:* (N₃N)Zr[C(PhPh)=NCH₂Ph] (**2**). A 6 mL benzene solution of (N₃N)ZrPPh (86 mg, 0.154 mmol) was cooled to ca. 5 °C, and a 2 mL benzene solution of PhCH₂N≡C (18 mg, 0.154 mmol) was added. The resultant orange solution was then frozen, and the benzene removed by lyophilization to give a pale orange powder (98 mg, 0.144 mmol, 94%). ¹H (C₆D₆, 500.1 MHz): δ 7.750 (t, C₆H₆, 2 H), 7.554 (d, C₆H₆, 2 H), 7.202 (t, C₆H₆, 2 H), 7.102 (m, C₆H₆, 4 H), 6.038 (d, PH, J_{PH} = 259 Hz), 4.903 (s, CH₂, 2 H), 3.287 (br s, CH₂, 6 H), 2.484 (s, CH₂, 6 H), 0.090 (s, CH₃, 27 H). ¹³C (C₆D₆, 125.8 MHz): δ 263.21 (d, C=N, J_{PC} = 99.5 Hz), 137.93 (s, Ph), 135.75 (d, Ph, J_{PC} = 16.6 Hz), 129.34 (s, Ph), 129.01 (s, Ph), 128.68 (s, Ph), 128.66 (d, Ph, J_{PC} = 6.9 Hz), 128.48 (s, Ph), 126.56 (s, Ph), 63.27 (d, CH₂, J_{PC} = 19.4 Hz), 61.21 (s, CH₂), 47.45 (s, CH₂), 2.19 (s, CH₃). ³¹P{¹H} (C₆D₆, 202.4 MHz): –39.98 (s). IR (KBr, Nujol): 2280 s (ν_{PH}), 1705 s (ν_{CN}) cm^{–1}. (N₃N)Zr[N(CH₂Ph)C(H)=PPh] (**3**). A 3 mL benzene solution of (N₃N)ZrPPh (173 mg, 0.309 mmol) and PhCH₂N≡C (36 mg, 0.309 mmol) was heated to 90 °C for 3 h. The orange solution was then frozen, and the benzene removed by lyophilization to give an orange powder, which was extracted into ca. 3 mL Et₂O. The orange solution was filtered then cooled to –30 °C to yield pale yellow crystals in several crops (163 mg, 0.241 mmol, 78%). ¹H (C₆D₆, 500.1 MHz): 10.291 (br s, CH, 1 H), 7.836 (br, C₆H₆, 2 H), 7.543 (br, C₆H₆, 2 H), 7.260 (t, C₆H₆, 2 H), 7.092 (m, C₆H₆, 2 H), 7.045 (m, C₆H₆, 2 H), 5.207 (s, CH₂, 2 H), 3.204 (s, CH₂, 6 H), 2.278 (s, CH₂, 6 H), 0.170 (s, CH₃, 27 H). ¹³C (C₆D₆, 125.8 MHz): δ 193.12 (d, C=N, J_{PC} = 55.6 Hz), 138.76 (s, Ph), 133.35 (s, Ph), 132.88 (d, Ph, J_{PC} = 16.6 Hz), 128.46 (s, Ph), 128.28 (s, Ph), 126.71 (s, Ph), 126.37 (s, Ph), 65.73 (s, CH₂), 64.46 (s, CH₂), 46.78 (s, CH₂), 1.38 (s, CH₃), one phenyl carbon resonance was not observed, presumably obscured by solvent. ³¹P{¹H} (C₆D₆, 202.4 MHz): 91.85 (s). Anal. Calcd for C₂₉H₅₂N₅PSi₃Hf: C, 51.43; H, 7.74; N, 10.34. Found: C, 51.54; H, 7.44; N, 10.63.
- ⁸ *Crystal data:* For **3**, C₂₉H₅₂N₅PSi₃Zr, *M* = 677.22, monoclinic, *P*2₁/c, *a* = 16.5616(8), *b* = 10.9871(5), *c* = 21.170(1) Å, β = 110.889(1)°, *Z* = 4, *V* = 3598.9(3) Å³, *T* = 125(2) K, μ(Mo–Kα) = 0.475 mm^{–1}. Of 49757 total reflections (pale yellow block, 1.35 = θ = 30.45°), 10366 were independent (*R*_{int} = 3.51%). The structure was solved using direct methods and standard difference map techniques and refined by full-matrix least-squares procedures on *F*². All non-hydrogen atoms were refined anisotropically. Hydrogen atoms on carbon were included in calculated positions and were refined using a riding model except the hydrogen atom on the phosphalkene carbon, H(16), which was located in the Fourier difference map and refined. *R*(*F*) = 2.89%, *R*(*wF*) = 6.78%, GoF = 1.033. CCDC reference number XXXX.

- 1 F. Mathey, *Phosphorus-Carbon Heterocyclic Chemistry: The Rise of a New Domain*; Pergamon: Amsterdam, 2001; E. Niecke, A. Fuchs, F. Baumeister, M. Nieger and W. W. Schoeller, *Angew. Chem., Intl. Ed.* **1995**, *34*, 555–557; J. I. Bates and D. P. Gates, *J. Am. Chem. Soc.*, **2006**, *128*, 15998–15999; D. Martin, F. S. Tham, A. Baccareddo and G. Bertrand, *Chem.–Eur. J.*, **2006**, *12*, 8444–8450.
- 2 F. Ozawa and M. Yoshifuji, *Dalton Trans.*, **2006**, *42*, 4987–4995; M. Yoshifuji, *Pure App. Chem.*, **2005**, *77*, 2011–2020.
- 3 I. Manners, *Angew. Chem., Intl. Ed.*, **2007**, *46*, 1565–1568; T. Baumgartner and R. Reau, *Chem. Rev.*, **2006**, *106*, 4681–4727; K. J. T. Noonan and D. P. Gates, *Angew. Chem., Intl. Ed.* **2006**, *45*, 7271–7274; V. A. Wright, B. O. Patrick, C. Schneider and D. P. Gates, *J.*

- Am. Chem. Soc.*, **2006**, *128*, 8836–8844; R. C. Smith, J. D. Protasiewicz, *J. Am. Chem. Soc.*, **2004**, *126*, 2268–2269.
- 4 H. W. Kroto and J. F. Nixon, *ACS Symp. Ser.*, **1981**, *171*, 383–390; L. N. Markovski and V. D. Romanenko, *Tetrahedron*, **1989**, *45*, 6019–6090; R. Appel *Multiple Bonds and Low Coordination in Phosphorus Chemistry*; M. Regitz, and O. J. Scherer, Eds.; Thieme Verlag: Stuttgart, Germany, 1990, p. 157; F. Mathey, *Acc. Chem. Res.*, **1992**, *25*, 90–96.
- 5 K. B. Dillon, F. Mathey and J. F. Nixon, *Phosphorus: The Carbon Copy*; John Wiley & Sons: New York, 1998, p. 88; P. P. Power, *Chem. Rev.*, **1999**, *99*, 3463–3504; M. Yoshifuji, *Phosphorus, Sulfur Silicon Relat. Elem.*, **2002**, *177*, 1827–1831; M. Yam, C.-W. Tsang and D. P. Gates, *Inorg. Chem.*, **2004**, *43*, 3719–3723.
- 6 R. Waterman, *Organometallics*, **2007**, *26*, 2492–2494; A. J. Roering, S. N. MacMillan, J. M. Tanski and R. Waterman, *submitted*.
- 7 D. M. Roddick, B. D. Santarsiero, J. E. Bercaw, *J. Am. Chem. Soc.*, **1985**, *107*, 4670–4678; Insertion of CO₂ into the Hf–P bond of Cp*₂HfH(PPh₂): G. A. Vaughan, G. L. Hillhouse, A. L. Rheingold, *Organometallics*, **1989**, *8*, 1760–1765.
- 8 F. Lindenberg, J. Sieler and E. Hey-Hawkins, *Polyhedron*, **1996**, *15*, 1459–1471.
- 9 D. W. Stephan, *Angew. Chem. Int. Ed.* **2000**, *39*, 314–329 and references therein.
- 10 Photochemical *E/Z* isomerization energies have been measured recently for *m*-terphenyl-supported phosphalkenes: Gudimetla, V. B.; Rheingold, A. L.; Payton, J. L.; Peng, H.-L.; Simpson, M. C.; Protasiewicz, J. D. *Inorg. Chem.* **2006**, *45*, 4895–4901.
- 11 K. Paasch, M. Nieger and E. Niecke, *Angew. Chem., Intl. Ed.* **1995**, *34*, 2369–2370.
- 12 J. F. Nixon, *Chem. Rev.*, **1988**, *88*, 1327–1362.
- 13 R. Appel, J. Peters, and R. Schmitz, *Z. Anorg. Allg. Chem.*, **1981**, *475*, 18–26.

Reaction of $(N_3N)ZrPPh$ (**1**) with $PhCH_2N\equiv C$ gave the 1,1-insertion product
 205 $(N_3N)Zr[C(PHPh)=NCH_2Ph]$ (**2**), which thermally rearranges to the structurally characterized
 phosphalkene-containing complex, $(N_3N)Zr[N(CH_2Ph)C(H)=PPh]$ (**3**), with perfect atom economy.

Graphical Contents Entry