

2015

Increasing Advanced Directive Utilization in Primary Care

Alejandro Castro
University of Vermont

Follow this and additional works at: <https://scholarworks.uvm.edu/fmclerk>



Part of the [Medical Education Commons](#), and the [Primary Care Commons](#)

Recommended Citation

Castro, Alejandro, "Increasing Advanced Directive Utilization in Primary Care" (2015). *Family Medicine Clerkship Student Projects*. 42.
<https://scholarworks.uvm.edu/fmclerk/42>

This Book is brought to you for free and open access by the Larner College of Medicine at ScholarWorks @ UVM. It has been accepted for inclusion in Family Medicine Clerkship Student Projects by an authorized administrator of ScholarWorks @ UVM. For more information, please contact donna.omalley@uvm.edu.

1. Increasing Advanced Directive Utilization in Primary Care

Hinesburg, VT

Alejandro Castro

Family Medicine Rotation 6 Dec 2014 – Jan 2015

Dr. James Ulager

2a. Problem Identification and Need

Advance directives are a vital tool ensuring patient's wishes regarding end-of-life care are met ^(1,2)

- End-of-life discussions with physicians are associated with fewer aggressive interventions and greater quality of life metrics
- Increased utilization of advance directives by Vermonters could ensure the wishes of patients are met
- A PRISM Analysis of all Patients ≥ 65 years of age at Hinesburg Family Practice with appointments in December 2014 showed 42.8% have an advance directive on file.
- Further analysis of these patients with specific billing codes on in the problem list (COPD, CHF, ACS/CVD, malignancy, CKD) showed 49.3% have an advance directive on file
- These rates are greater than one large study of CHF patients but still represent tremendous opportunity

2b. Problem Identification and Need

- Additional data pertaining to Vermonter's wishes regarding end-of-life care³
 - 80% of Vermonters indicate they would like to die at home yet only 23% achieve this goal
 - The rate of Vermonters able to die at home improves from 23% to 76% for those enrolled in hospice

3. Public Health Costs

- Palliative care consultations for MICU patients was associated with greater utilization of DNR orders and reduced time to death⁴
- Vermont ranks near the 10th percentile of state enrollment in hospice care in the last 6 months of life (20.2% of Medicare decedents)⁵
 - Vermont Medicare spending last 2 years of life: \$41,514
 - Vermont hospice care spending: \$1382
- Although the principal purpose of improved end-of-life care is *not* cost containment, greater utilization of advance directives, DNR orders as well as palliative and hospice services has the potential for significant cost savings

4a. Community Perspectives

- [Name Withheld], LICSW - Social Worker, Community Health Team
 - Confirmed the underutilization of advanced directives and discussions regarding end-of-life wishes
 - Highlighted the ongoing nature of end-of-life care discussions involving a back and forth with providers and changing goals over time
 - Recommended interventions that incorporate family members and a continuous evaluation of what a patient's beliefs are and what their wishes would be

4a. Community Perspectives

- Dr. [Name Withheld] M.D - researcher in palliative care and end-of-life communication in the ICU
 - Also highlighted the underutilization of advance directives as well as palliative and hospice services
 - Indicated there is lack of awareness within the community and wide variance among providers in the timing and responsibility of completing an advance directive
 - Highlighted the importance of timely conversations with dying patients regarding their prognosis as a means of closing the disconnect between patient preferences and what actually occurs nearing the end of life

5. Intervention and Methodology

- Identify utilization of advance directives within Hinesburg Family Practice
- Identify demographics with particular life-limiting illnesses that are most in need of advance directives
- Develop accessible literature that can be provided more regularly to the targeted demographics

6. Results

- An effective pamphlet was developed that distilled the salient information from multiple sources into a single piece of literature for:
 - Understanding what an advance directive is
 - How to complete an advance directive
 - Where and with whom advance directives are submitted
 - Location for accessing additional information
- This pamphlet is provided frequently and cheaply to specific patient demographics

7. Evaluation

- Evaluation will involve again pulling data from PRISM for the identified demographics on slide 2a
- Over time there should be an statistically significant increase in advance directives on file for targeted patients

8. Future Recommendations

- When considering implementing new guidelines, interventions, or protocols, administrators must balance the value of an intervention and the cost of overwhelming providers
- PRISM can identify targeted cohorts of patients and limit burdens on providers
- Strong consideration should be given to include a prompt within the “COPD Action Plan” for patients below certain physiologic thresholds that indicates a provider has discussed advance directives or one is already on file
 - This would be narrow in scope so as not to overwhelm providers
 - An important demographic with a valuable intervention would be targeted

9. References

1. Wright AA, Zhang B, Ray A, et al. Associations Between End-of-Life Discussions, Patient Mental Health, Medical Care Near Death, and Caregiver Bereavement Adjustment. *JAMA*. 2008;300(14):1665-1673. doi:10.1001/jama.300.14.1665.
2. Advance Directives Among Hospitalized Patients With Heart Failure, *JACC: Heart Failure*, Volume null, Issue null, Page null, Javed Butler, Zachary Binney, Andreas Kalogeropoulos, Melissa Owen, Carolyn Clevenger, Debbie Gunter, Vasiliki Georgiopoulou, Tammie Quest
3. No. 60. An act relating to hospice and palliative care, <http://www.leg.state.vt.us/DOCS/2012/ACTS/ACT060.PDF>
4. Palliative & Supportive CarePB - Cambridge Journals OnlineAU - Lustbader,DanaAU - Pekmezaris,ReneeAU - Frankenthaler,MichaelAU - Walia,RajniAU - Smith,FrederickAU - Hussain,ErfanAU - Napolitano,BarbaraAU - Lesser,MartinTI - Palliative medicine consultation impacts DNR designation and length of stay for terminal medical MICU patientsSN - 1478-9523PY - 2011VL - 9IS - 04SP - 401EP - 406M3 - 10.1017/S1478951511000423UR - <http://dx.doi.org/10.1017/S1478951511000423>ER -
5. Percent of decedents enrolled in hospice during the last six months of life, The Dartmouth Atlas of Health Care