

University of Vermont

UVM ScholarWorks

College of Nursing and Health Sciences Doctor
of Nursing Practice (DNP) Project Publications

College of Nursing and Health Sciences

2020

Improving Surveillance of Hepatitis C Infections Among Patients Receiving Medication Treatment for Opioid Use Disorder

Jacqueline Bray
University of Vermont

Follow this and additional works at: <https://scholarworks.uvm.edu/cnhsdnp>



Part of the [Nursing Commons](#)

Recommended Citation

Bray, Jacqueline, "Improving Surveillance of Hepatitis C Infections Among Patients Receiving Medication Treatment for Opioid Use Disorder" (2020). *College of Nursing and Health Sciences Doctor of Nursing Practice (DNP) Project Publications*. 40.

<https://scholarworks.uvm.edu/cnhsdnp/40>

This Project is brought to you for free and open access by the College of Nursing and Health Sciences at UVM ScholarWorks. It has been accepted for inclusion in College of Nursing and Health Sciences Doctor of Nursing Practice (DNP) Project Publications by an authorized administrator of UVM ScholarWorks. For more information, please contact scholarworks@uvm.edu.

Improving Surveillance of Hepatitis C Infections Among Patients Receiving Medication Treatment for Opioid Use Disorder

Jacqueline Bray, BA, RN, DNP-c; Academic Advisor: Deborah Wachtel DNP, MPH, APRN; Site Advisor: Robin Sherman APRN-BC

Background

- Abuse of opioids has shifted from oral to injection use, resulting in greater risk of infectious disease transmission
- The number of new Hepatitis C virus (HCV) infections in the US has tripled
- Treating acute infections of HCV vs. managing chronic Hepatitis C is expected to result in \$12 billion in total cumulative savings to Medicaid by 2022
- Despite engagement in treatment, individuals with opioid use disorder (OUD) remain at risk for contracting and transmitting infectious disease, yet many mOUD programs screen patients upon admission only.



Purpose

- Implement a HCV re-screening protocol in a mOUD program and determine sustainability of protocol
- Secondarily, determine if point of care finger stick tests would be a preferred modality to use for screening


Methods

- Patients with an unknown or negative HCV antibody test result upon admission were offered to complete a risk survey
 - Patients were given a list of risk factors for HCV, and asked to answer if they applied YES/NO
 - IF YES, patients were asked if they would like be re-screened for HCV
- If interested in rescreening, patients were offered an onsite venipuncture
- Patients who declined rescreening, were asked if they would re-consider if the testing modality was a finger stick
- Patients with a positive HCV RNA viral load received a referral to UVMHC Infectious Disease
- Nursing staff, medical providers, and administrative leaders provided feedback via an online survey

Results



- Survey Questions
- The risk survey was easy to administer to patients.
 - Administering the risk survey at the dosing window did not interrupt work flow.
 - Administering the risk survey at the dosing window did not interfere with responsibilities of the dosing nurse.
 - Entering and updating flags in the electronic health system to facilitate surveillance of HCV is feasible to add to nursing work flow.
 - I am comfortable performing venipuncture on patients who wish to be re-screened at the clinic.
 - I would be comfortable performing a finger stick at the dosing window to collect a sample of blood to re-screen patients for HCV if proper training were provided.
 - Having patients complete the risk survey at the window presents a brief opportunity to educate patients about the risk factors for Hepatitis C and the importance of routine screening.
 - Dispensing HCV medication with a patient's daily dose is easy and is an effective strategy to promote medication adherence.



- Survey Questions
- It is important to re-screen patients who had a negative HCV antibody result upon admission to the Chittenden Clinic.
 - It is important to offer re-screening for HCV antibody to patients at the Chittenden Clinic 6 months after the initial negative result.
 - It is important to offer re-screening for HCV antibody to patients at the Chittenden Clinic 1 year after the initial negative result.
 - It is within the Chittenden Clinic's scope of practice and treatment program to offer health care services such as screening, diagnosing, and care coordination for patients at risk for and/or who have an infectious disease.
 - Re-screening for infectious disease, is the responsibility of the patient's primary care provider or public health service team, and not the responsibility of the Chittenden Clinic.
 - Point of care testing for HCV is a service that should be offered to patients at the Chittenden Clinic.
 - It is important for Chittenden Clinic providers to be able to prescribe medications to treat Hepatitis C.
 - The Chittenden Clinic offers a unique environment in which the identification and treatment of Hepatitis C can be managed effectively.
 - It is the role and responsibility of the Chittenden Clinic care team to educate patients on the risk of contracting Hepatitis C.
 - The Chittenden Clinic should prioritize improving the surveillance of infectious disease among its patients.

Discussion

- A brief questionnaire administered to patients regarding risk factors for HCV can help facilitate targeted screening efforts
- Uptake of re-screening may improve if less invasive screening modalities were used
- Nursing staff agreed that re-screening surveys were easy to implement, and introduced opportunities to talk with patients about their risk factors for contracting HCV, and review the importance of routine screening
- Stakeholders agreed that routine surveillance of HCV is important, and fits within the the scope of practice of mOUD treatment

Implications for Future Practice

- Risk questionnaires could be distributed to patients receiving medication for OUD in primary care settings
- Surveillance services offered in mOUD treatment programs may qualify for enhanced-rate reimbursement
- In addition to on-site identification of HCV, treatment of HCV in a mOUD setting is cost effective and patient centered

Conclusion

- Integration of HCV surveillance services at mOUD treatment programs provides a sustainable approach to reduce the public health burden of infectious disease



THE UNIVERSITY OF VERMONT
COLLEGE OF NURSING
AND HEALTH SCIENCES

The University of Vermont