

2016

Small Grain Disease and Insect Pest Scouting Report

Heather Darby

University of Vermont, heather.darby@uvm.edu

Ann Hazelrigg

University of Vermont

Erica Cummings

University of Vermont

Gabriella Maia

University of Vermont

Follow this and additional works at: <https://scholarworks.uvm.edu/nwcsp>



Part of the [Agricultural Economics Commons](#)

Recommended Citation

Darby, Heather; Hazelrigg, Ann; Cummings, Erica; and Maia, Gabriella, "Small Grain Disease and Insect Pest Scouting Report" (2016). *Northwest Crops & Soils Program*. 102.

<https://scholarworks.uvm.edu/nwcsp/102>

This Report is brought to you for free and open access by the UVM Extension at ScholarWorks @ UVM. It has been accepted for inclusion in Northwest Crops & Soils Program by an authorized administrator of ScholarWorks @ UVM. For more information, please contact donna.omalley@uvm.edu.



2016 Small Grain Disease and Insect Pest Scouting Report

Dr. Heather Darby, Extension Agronomist
Dr. Ann Hazelrigg, Director, Plant Diagnostic Clinic
Erica Cummings, Crops and Soils Coordinator
Gabriella Maia, Plant Diagnostic Clinic Research Assistant
(802) 524-6501

Visit us on the web at <http://www.uvm.edu/extension/cropsoil/>

INTRODUCTION

A survey of small grain diseases and insect pests were conducted in Vermont, New York, and Massachusetts during the 2016 growing season. Pests were scouted on six Vermont farm locations in the towns of Alburgh, North Troy, Shelburne, Bridport, Glover, and Berlin, as well as in Essex, New York and Northfield, Massachusetts.

Winter and spring wheat (heirloom and commercially available) as well as spring barley were scouted between spike emergence and flowering, and again at the soft dough growth stage. Disease and insect samples were taken and identified with assistance from the University of Vermont (UVM) Plant Diagnostic Clinic.

DISEASES IDENTIFIED

The overall warm and dry growing conditions throughout much of the season resulted in relatively low levels of foliar diseases. The foliar and head diseases identified during the 2016 growing season are described below:

Foliar Diseases

Foliar diseases reduce photosynthetic leaf area, use nutrients, and increase respiration and transpiration within the infected plant tissues. A diseased plant typically exhibits reduced vigor, growth, and seed fill. The earlier the occurrence, the greater degree of infection, and the longer duration of conditions favorable for disease development, the yield loss will be greater. The following foliar diseases were identified during scouting visits.



Image 1. Powdery mildew on spring wheat.

Powdery mildew (*Erysiphe graminis f. sp. Tritici*) was identified on the leaves and stems of winter wheat in

Northfield, MA, on the leaves of winter wheat in Shelburne, and on the leaves of winter and spring wheat and spring barley in Alburgh, VT. Powdery mildew is relatively easy to identify, the fungus produces whitish-gray cottony growths on the upper leaf surface or stem of the infected plant (Image 1). Infection primarily occurs on the lower leaves and stem sections of the plant.



Image 2. Tan spot infected leaves (*Pyrenophora tritici-repentis*).

Stagonospora spreads, the yellowing increases and forms lens-shaped blotches on the leaf that eventually turns red-brown. As *Stagonospora* progresses, the lesions develop an ashen gray-brown center containing brown specks (but without the distinct yellow border typical of tan spot lesions).

Leaf rust (*Puccinia recondite*) was observed at only two sites, Alburgh, VT and Northfield, MA. Leaf rust is the most common of the rust pathogens. It is characterized by round rusty-red/orange masses of spores on the leaf surface (Image 4). **Striped rust (*Puccinia striiformis*)** was also identified on winter wheat at the Northfield, MA location.

Leaf spots such as **Tan spot (*Pyrenophora tritici-repentis*)**, **Septoria tritici blotch (STB) (*Zymoseptoria tritici*)** and **Stagonospora leaf and glume blotch (*Stagonospora nodorum*)** were identified at all of the on-farm sites in Vermont, Massachusetts, and New York, and on all grain types.

The Tan Spot fungus produces elongated asymmetrical spots (1/8 to 1/2-inch long and 1/16 to 1/18-inch wide) (Image 2). Here, a tiny, dark spot forms (best observed by holding the leaf up to the light). The spot enlarges into a tan lesion, surrounded with a narrow to broad yellow border to produce an “eyespot” type of symptom, characteristic of this disease.

Septoria tritici blotch (STB) (*Zymoseptoria tritici*) and *Stagonospora* leaf and glume blotch (*Stagonospora nodorum*) look very similar and are often confused with one another (Image 3). Both start out as yellow spots. However, as STB spreads, irregular brown lesions form along leaf veins giving the appearance of stripes. In the middle of these lesions, dark brown spore masses (pycnidia) form that can be seen with the naked eye, making a distinguishing characteristic between these two diseases. In contrast, as



Image 3. Left leaf infected with *Septoria tritici* blotch (STB) (*Zymoseptoria tritici*) and the right with *Stagonospora* leaf blotch (*Stagonospora nodorum*).



Image 4. Leaf rust (*Puccinia recondite*) on winter wheat.



Image 5. Genetic resistance response in winter wheat.

Interestingly a genetic resistance response to foliar diseases was observed on wheat at all sites (Image 5).

Grain Head Diseases

There are two primary grain head diseases found in the Northeast, **Loose smut** (*Ustilago tritici*) and **Fusarium head blight** (FHB) (*Fusarium graminearum*). Loose smut was identified at the Alburgh site on spring and winter wheat and spring barley, and on winter wheat at the Bridport and Shelburne, VT locations. The loose smut fungus is carried as dormant mycelium within healthy-looking seed and is spread by planting infected seed. A smut-infected seed and plant cannot be distinguished from an uninfected one until the head starts to emerge. The disease is most obvious just after the time of heading by the characteristic dusty black appearance of diseased heads (Image 6). The spores are dispersed by the wind during wheat flowering and can infect healthy plants.



Image 6. Loose smut (*Ustilago tritici*) infected wheat head.

If you find heads with loose smut in your fields, you should NOT save the seed for future planting. Loose smut is not considered a human health risk, but planting infected seed will exponentially increase diseased seed and result in yield losses.

The pathogen of most concern among grain growers is *Fusarium* head blight (FHB). It is predominantly caused by the species, *Fusarium graminearum*. This disease is very destructive and causes yield losses, low test weights, low seed germination, and contamination of grain with the mycotoxin, a vomitoxin, called deoxynivalenol (DON). The spores are usually transported by air currents and can infect plants at flowering through grain fill. Spores can also overwinter on grain stubble. A telltale sign of FHB infection is the premature bleaching of grain heads. Another symptom is a pink or orange colored mold at the base of the spikelet. Additionally, once the grains are harvested, infected kernels will be pink, white, chalky and/or shriveled. *Fusarium* can



Image 7. Incomplete grain fill of winter wheat.

pose a health risk to both humans and livestock. Consumption of contaminated grains at DON levels of greater than 1 ppm in humans and 10 ppm for certain livestock can cause illness; therefore, it is critically important to test grain for DON. More information on DON testing can be found at:

<http://www.uvm.edu/extension/cropsoil/cereal-grain-testing-lab>.

All on-farm locations were scouted at soft dough to assess FHB severity using the North Dakota State University visual scale. The warm and dry conditions during the 2016 growing season resulted in minimal observations of FHB infection at any of the on-farm sites and record low DON levels.

Interestingly, while scouting for FHB infection we observed incomplete grain fill of predominantly winter wheat, at several of the on-farm locations, which resulted in lower than expected yields (Image 7). This might be due to drought like conditions during grain fill.

Managing Grain Diseases

It is important to remember, we do not know directly how foliar diseases affect yields. Although we have identified these issues in the field, it is not clear as to how, or at what severity, they impact yield and quality.

In our cool, moist climate, practices that are critical to managing the multitude of diseases that impact small grains include: planting clean seed, rotating crops, and improving air flow. We highly recommend buying “certified” seed when possible. Certified seed guarantees that the seed meets or exceeds a strict set of quality control standards. Weed management is important, especially in spring grains to improve airflow and assist with keeping the plants as dry as possible. Spores from many of the fungal diseases can survive in the soil or plant debris for several years waiting for their host plant and/or ideal conditions. Therefore, crop rotation and healthy soil is critical to minimizing diseases present during grain production. Conventional growers may purchase fungicide-treated seed to help mitigate some of the disease issues. There are also several commercial pesticides available as a last resort to control extreme outbreaks.

INSECT PESTS IDENTIFIED

Insect pests were more predominate this season most likely due to the drought like conditions during the growing season (Table 1). Thrips were identified at all scouting locations, however overall damage was minimal. Cereal leaf beetle damage was recorded in Alburgh, Berlin, North Troy, and Glover, VT, as well as in Northfield, MA and Essex, NY. The most severe cereal leaf beetle damage was measured in Northfield, MA on spring wheat. Interestingly, there were two varieties of spring wheat planted in this field, one had 85% damage and the other, only 5%, indicating that variety selection is important. Plant injury by the Brown Wheat Mite was identified at the Alburgh, North Troy, and Glover, VT scouted fields. Other insect pests

identified include; wireworm (Alburgh, North Troy, and Glover, VT), aphids (Bridport, VT and Northfield, MA), slugs (Berlin, VT and Glover, VT), and Mexican bean beetle (Glover, VT).

Table 1. Insect pests identified at the on-farm scouting locations

Location	Grain type	Cereal Leaf Beetle	Thrips	Brown Wheat Mite	Wire worm	Slugs	Mexican Bean Beetle	Aphids
Alburgh, VT	Spring Wheat	X	X	X	X			
Alburgh, VT	Winter Wheat	X	X	X	X			
Berlin, VT	HRWW	X	X			X		
Bridport, VT	HRWW		X					X
Bridport, VT	SWWW		X					
Essex, NY	Spring Barley	X	X					
Glover, VT	HRSW	X	X	X		X	X	
North Troy, VT	HRSW	X	X	X				
Northfeild, MA	HRWW	X	X					
Northfeild, MA	HRWW	X	X					
Northfeild, MA	SWWW	X	X					X
Northfeild, MA	HRSW	X	X					
Northfeild, MA	HRSW	X	X					
Shelburne, VT	HRWW		X					
Shelburne, VT	SWWW		X					

With the exception of cereal leaf beetle damage on spring wheat in Northfield, MA, the insect pests observed at the on-farm locations did not appear to pose a major threat to the grain crops.

SCOUTING TIPS

Start scouting grains for foliar diseases and insect pests at the flag leaf growth stage, and be on the lookout for grain head diseases starting at spike emergence. Rogue out any smutted heads and do not save seed where loose smut is present. Keep an eye out for premature bleaching of grain heads and salmon colored fungus on spikelets, but remember, just because you have the fungus, does not necessarily mean you have the mycotoxin and vice versa—so be sure to test your grains for DON. After harvest, a grain sample was taken from each of the on-farm scouting sites and analyzed for DON concentration. All were under 1ppm.



This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2104-70006-22577. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture.

United States
Department of
Agriculture

National Institute
of Food and
Agriculture

UVM Extension helps individuals and communities put research-based knowledge to work. Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. University of Vermont Extension, Burlington, Vermont. University of Vermont Extension, and U.S. Department of Agriculture, cooperating, offer education and employment to everyone without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or familial status.

