

University of Vermont

UVM ScholarWorks

College of Nursing and Health Sciences Doctor
of Nursing Practice (DNP) Project Publications

College of Nursing and Health Sciences

2022

ERAS: Improving Surgical Experience in Traumatic Hip Fracture Surgery

Kristen Knight DNP(c)
University of Vermont

Marcia Bosek, DNSc
The University of Vermont

Follow this and additional works at: <https://scholarworks.uvm.edu/cnhsdnp>



Part of the [Nursing Commons](#)

Recommended Citation

Knight, Kristen DNP(c) and Bosek, DNSc, Marcia, "ERAS: Improving Surgical Experience in Traumatic Hip Fracture Surgery" (2022). *College of Nursing and Health Sciences Doctor of Nursing Practice (DNP) Project Publications*. 102.

<https://scholarworks.uvm.edu/cnhsdnp/102>

This Project is brought to you for free and open access by the College of Nursing and Health Sciences at UVM ScholarWorks. It has been accepted for inclusion in College of Nursing and Health Sciences Doctor of Nursing Practice (DNP) Project Publications by an authorized administrator of UVM ScholarWorks. For more information, please contact schwyrks@uvm.edu.



ERAS: Improving Surgical Experience in Traumatic Hip Fracture Surgery

Kristen Knight, DNP(c), RN
 Project Advisor: Marcia Bosek, DNSc, RN
 Site Mentor: Kathleen Gouldy, MSN, APRN
 The University of Vermont



THE UNIVERSITY OF VERMONT
 COLLEGE OF NURSING
 AND HEALTH SCIENCES

Background

- Hip fractures are associated with significant morbidity and mortality, loss of independence and financial burden.¹
- Enhanced Recovery After Surgery (ERAS) pathways are utilized to improve postoperative outcomes for treatment of hip fracture without increasing rates of complication or mortality.²

Purpose and Aims

Purpose: To improve perioperative care for traumatic hip fracture patients by utilizing ERAS pathway.

Aim I. Create an ERAS Orthopedic Pathway (ERAS-OP) for traumatic hip fracture.

Aim II a. Evaluate the consistency of implementation of the post-operative Phase of ERAS-OP.

Aim II b. Evaluate post-operative ERAS-OP outcomes: length of stay (LOS), cost of care & opioid administration.

Methodology

- This quality improvement project takes place at an urban Midwest trauma II center.
- 128 hip fracture patients met the eligibility criteria (66 ERAS, 62 control).
- Data were collected on patients treated for traumatic hip fracture with post-operative ERAS-OP (ERAS: June–September 2021), compared to a retrospective cohort of controls managed without ERAS-OP (Control June–September 2020).

Aim I: Utilizing Evidence Based Practices to Create ERAS Orthopedic Pathway

- A review of the literature revealed evidence based practices specific to hip fracture along with basic ERAS components.

Basic ERAS Components	Hip Fracture Specific Components
<ul style="list-style-type: none"> Pre-operative education Oral multimodal analgesia Fluid optimization Early mobilization Early oral intake Scheduled alternatives to opioids, nausea and vomiting control and supported discharge. Pre-operative fasting and carbohydrate treatment Prevention of Post-Operative Nausea and Vomiting 	<ul style="list-style-type: none"> Time to surgery Regional anesthesia Delirium Prevention Early advanced care planning

Aim II a: Consistency of Implementation: Post-Operative Protocols

- Time to Nutrition: Full Liquid, Soft or Solid Diet type and diet or snack percentage consumed (> 0%) documented within 13 hours of exiting the OR.
- Time to Ambulation: Documented therapeutic ambulation within 7 hours (13 hours total joints) of leaving the OR.
- Sustained Ambulation Pass: Documented therapeutic ambulation 3 times within 24 hours of leaving the OR.
- Scheduled Alternative to Opioids (ALTO): Verified order of non-opioid medication with scheduled frequency (not prn) for administration on inpatient units.

Aim II b: Post-Operative Outcomes


- To compare outcomes pre and post implementation, metrics data (as defined below) were collected from June 2020 through September 2020 (control) and from June 2021 through September 2021 (ERAS).

- Length of Stay:** Length of stay from out of operating room event time to discharge.
- Morphine Milligram Equivalent (MME) administered:** MME administered for all oral and IV pushes from out-of-PACU until discharge.
- Total Cost:** Costs accrued during the patient's entire length of stay.

Results

Aim I: Utilizing Evidence Based Practices to Create ERAS Orthopedic Pathway

- The ERAS-OP was approved by the ERAS systems coordinator at the specified trauma II hospital (to see ERAS-OP scan QR)
- Only the post-operative ERAS-OP interventions were in effect as of 6/1/2021.
- The remainder of ERAS-OP will be implemented by 5/1/2022.





Aim II a : Consistency of Implementation: Post-Operative Protocols (n= 128)



- 42% Time to Nutrition Pass
- 33% Time to Ambulation
- 14% Sustained Ambulation
- 80% Scheduled Alternatives to Opioids

Aim II b: Post-Operative Implementation



2020 (n=62) vs 2021 (n=66) length of stay from time out of operating room to time of discharge (p=.94)



2020 (n=62) vs 2021 (n=66) morphine milligram equivalents (MME) administered from out-of-PACU until discharge (p=.09)



2020 (n=62) vs 2021 (n=66) costs accrued during the entire length of stay of the patient (p < .001)



Discussion

Aim I

- Two barriers to implementation were identified: NPO time to surgery and regional anesthesia administration.
- Further institutional investigation is warranted to identify the scope of these barriers.

Aim II

This cohort of traumatic hip fracture patients treated with post-operative ERAS-OP demonstrated:

- No significant change to length of stay (p=.94).
 - No reduction in LOS may be due to a shortage of beds available in skilled nursing facilities and rehab facilities, thus prolonging inpatient LOS.
- Inadequate evidence to support decrease of MME (p =.09).
 - Reduction in post-operative MME administration is likely the result of providing scheduled alternative to opioid medications and encouraging early ambulation after surgery.
- Improved outcomes related to:
 - Lower total cost of inpatient care (p < .0001).
 - Reduction in inpatient cost may be the result of the ERAS-OP helping to minimize post-operative complications and costly medication administration.

Conclusions

- This cohort of traumatic hip fracture patients treated with post-operative ERAS-OP demonstrated a decrease in cost of care.
- The findings from this QI project support the benefit of ERAS for use with traumatic hip fractures.
- Further evaluation will be needed once the entire ERAS-OP is implemented in 2022.

Future Considerations

- Further research is needed to better understand:
 - Barriers to NPO guidelines.
 - Implementation of regional anesthesia in the Emergency Department.
 - Barriers to provider, nursing and patient compliance with protocol.

References

- Kang, Y., Liu, J., Chen, H., Ding, W., Chen, J., Zhao, B., & Yin, X. (2019). Enhanced recovery after surgery (ERAS) in elective intertrochanteric fracture patients result in reduced length of hospital stay (LOS) without compromising functional outcome. *Journal of Orthopedic Surgery and Research*, 14(1), 209-209. <https://doi.org/10.1186/s13018-019-1238-2>
- Schnell, S., Friedman, S. M., Mendelson, D. A., Bingham, K. W., & Kates, S. L. (2010). The 1-year mortality of patients treated in a hip fracture program for elders. *Geriatric Orthopedic Surgery & Rehabilitation*, 1(1), 6-14. <https://doi.org/10.1177/2151458510378105>