

2012

Enhancing Forages with Nutrient Dense Sprays 2012 Trials

Heather Darby

University of Vermont, heather.darby@uvm.edu

Susan Monahan

University of Vermont

Erica Cummings

University of Vermont

Hannah Harwood

University of Vermont

Rosalie Madden

University of Vermont

Follow this and additional works at: <https://scholarworks.uvm.edu/nwcsp>



Part of the [Agricultural Economics Commons](#)

Recommended Citation

Darby, Heather; Monahan, Susan; Cummings, Erica; Harwood, Hannah; and Madden, Rosalie, "Enhancing Forages with Nutrient Dense Sprays 2012 Trials" (2012). *Northwest Crops & Soils Program*. 247.

<https://scholarworks.uvm.edu/nwcsp/247>

This Report is brought to you for free and open access by the UVM Extension at ScholarWorks @ UVM. It has been accepted for inclusion in Northwest Crops & Soils Program by an authorized administrator of ScholarWorks @ UVM. For more information, please contact donna.omalley@uvm.edu.



Enhancing Forages with Nutrient Dense Sprays 2012 Trials



Dr. Heather Darby, UVM Extension Agronomist
Susan Monahan, Erica Cummings, Hannah Harwood and Rosalie Madden
UVM Extension Crops and Soils Technicians
802-524-6501

Visit us on the web: <http://www.uvm.edu/extension/cropsoil>

ENHANCING FORAGES WITH NUTRIENT DENSE SPRAYS, 2012 TRIALS

Dr. Heather Darby, University of Vermont Extension

Heather.Darby@uvm.edu

INTRODUCTION

The nutrient dense study was initiated at two locations in Vermont to test the efficacy of amending forages with foliar sprays. The nutrient spray program was developed by Advancing Eco-Agriculture and consisted of five foliar sprays for the Vermont farms in this study. The recommended spray program included applications of Rejuvenate in the early spring and late fall, and a combination of PhotoMag, Phosphorus, Potassium and MicroPak applied in the spring and after each cut of hay or graze (Table 1). This study was conducted based on farmer interest in enhancing nutrient density of forages through foliar sprays and was funded by the Lattner Foundation. Any reference to commercial products, trade names or brand names is for information only, and no endorsement or approval is intended.

Table 1. Information on Advancing Eco-Agriculture nutrient dense sprays.¹

| Spray | What is it? | What does it do? |
|------------|--|---|
| Rejuvenate | humic substance, carbohydrates, sea minerals | stimulates soil microbial life |
| PhotoMag | magnesium, sulfur, boron, cobalt | promotes chlorophyll and sugar production |
| Phosphorus | mined phosphate | improves photosynthesis and plant root vigor |
| Potassium | mined potassium sulfate | improves storability |
| MicroPak | boron, zinc, manganese, copper, cobalt, molybdenum, sulfur | enhances sugar translocation, root strength, and plant immunity |

¹Information gathered from the Advancing Eco-Agriculture website: growbetterfood.com.

MATERIALS AND METHODS

In 2012, forages were amended with nutrient dense sprays at two locations: Shelburne Farms in Shelburne, VT and Butterworks Farm in Westfield, VT. Both hayfields had been in native grass/legume mixture for many years. The nutrient recommendations from Advancing Eco-Agriculture are listed in Table 2. In order to understand what may cause a response, if any, we compared the recommended spray regime ('All') to individual components, as well as a control of water. The experimental design was a randomized complete block with four replications.

Table 2. Timing and amount of Nutrient Dense Sprays used.

| Timing | Recommendations (per acre) |
|--------------------|--|
| Early Spring | 3 tons compost, 20 lb. Borate (10%), and 5 lbs. Zinc sulfate, 2 gallons Rejuvenate |
| After Each Cut | 1 gallon PhotoMag, 1 gallon Phosphorus, 1 quart Potassium, 2 quarts MicroPak |
| Fall, post harvest | 6 quarts Rejuvenate, 2-3 tons compost |

Six by ten foot plots were established in existing hay fields. Harvest and spray dates for each location are presented in Table 3. Plots were harvested with a BCS sickle bar mower (Portland, OR), raked by hand, gathered and weighed on a platform scale. A subsample was dried at 40° C and weighed to determine dry matter. Oven dry samples were coarsely ground with a Wiley mill (Thomas Scientific, Swedesboro, NJ) and sent to Cumberland Valley Analytical Services, Inc. (Hagerstown, MD) for quality analysis. Results were analyzed with an analysis of variance in SAS (Cary, NC).

Table 3. Harvest and spray dates at each location.

| Treatment | Butterworks Farm | Shelburne Farms |
|----------------------|------------------|---------------------|
| Spray Rejuvenate | 18-Apr | 19-Apr |
| Spray All Treatments | 16-May | 24-Apr (B, Zn only) |
| 1 st Cut | 31-May | 17-May |
| Spray All Treatments | 12-Jun | 29-May |
| 2 nd Cut | 9-Jul | 21-Jun |
| Spray All Treatments | 18-Jul | 5-Jul |
| 3 rd Cut | 21-Aug | 27-Jul |
| Spray All Treatments | 28-Aug | 7-Aug |
| Spray Rejuvenate | 9-Oct | 9-Oct |

SILAGE QUALITY

Silage quality was analyzed by Cumberland Valley Analytical Forage Laboratory in Hagerstown, Maryland. Plot samples were dried, ground and analyzed for crude protein (CP), acid detergent fiber (ADF), neutral detergent fiber (NDF) and various other nutrients. The Nonstructural Carbohydrates (NSC) and Total Digestible Nutrients (TDN) were calculated from forage analysis data. Performance indices such as Net Energy for Lactation (NEL) were calculated to determine forage value. Mixtures of true proteins, composed of amino acids and nonprotein nitrogen make up the crude protein (CP) content of forages. The bulky characteristics of forage come from fiber. Forage feeding values are negatively associated with fiber since the less digestible portions of the plant are contained in the fiber fraction. The detergent fiber analysis system separates forages into two parts: cell contents, which include sugars, starches, proteins, non-protein nitrogen, fats and other highly digestible compounds; and the less digestible components found in the fiber fraction. The total fiber content of forage is contained in the neutral detergent fiber (NDF). Chemically, this fraction includes cellulose, hemicellulose and lignin. Recently, forage testing laboratories have begun to evaluate forages for NDF digestibility. Evaluation of forages and other feedstuffs for NDF digestibility is being conducted to aid prediction of feed energy content and animal performance. Research has demonstrated that lactating dairy cows will eat more dry matter and produce more milk when fed forages with optimum NDF digestibility. Forages with increased NDF digestibility (dNDF) will result in higher energy values, and perhaps more importantly, increased forage intakes. Forage NDF digestibility can range from 20 – 80%. The NSC or non-fiber carbohydrates (NFC) include starch, sugars and pectins.

LEAST SIGNIFICANT DIFFERENCE (LSD)

Variations in yield and quality can occur because of variations in genetics, soil, weather and other growing conditions. Statistical analysis makes it possible to determine whether a difference among varieties is real, or whether it might have occurred due to other variations in the field. At the bottom of each table, a LSD value is presented for each variable (i.e. yield). Least Significant differences (LSD's) at the 10% level of probability are shown. Where the difference between two treatments within a column is equal to or greater than the LSD value at the bottom of the column, you can be sure in 9 out of 10 chances that there is a real difference between the two varieties. Treatments that were not significantly lower in performance than the highest value in a particular column are indicated with an asterisk. In the example below, A is significantly different from C but not from B. The difference between A and B is equal to 1.5 which is less than the LSD value of 2.0. This means that these varieties did not differ in yield. The difference between A and C is equal to 3.0 which is greater than the LSD value

| Variety | Yield |
|---------|-------|
| A | 6.0 |
| B | 7.5* |
| C | 9.0* |

of 2.0. This means that the yields of these varieties were significantly different from one another. The asterisk indicates that B was not significantly lower than the top yielding variety.

LSD **2.0**

RESULTS AND DISCUSSION

Seasonal precipitation and temperature recorded at weather stations in close proximity to Westfield and Shelburne, VT are reported in Table 4. The temperature and precipitation in Westfield was close to the 30-year average. There were a total of 5530 GDD (growing degree days), 134 GDD above average. May, August and October were warmer than average in Westfield, with less rain in July and August. In Shelburne, monthly temperatures were above the 30-year average every month of the growing season. There were a total of 6488 GDD, 639 GDD above average. Warmer temperatures in Shelburne contribute to the earlier harvests of 2nd and 3rd cut hay.

Table 4. Seasonal weather data collected near Westfield and Shelburne, VT.

| Westfield* | April | May | June | July | August | Sept | Oct |
|-------------------------------|-------|------|------|------|--------|------|------|
| Average Temperature (F) | 41.8 | 56.7 | 63.0 | 67.9 | 68.1 | 56.9 | 48.8 |
| Departure from Normal | -0.9 | 1.9 | -0.8 | -0.1 | 2.0 | -0.6 | 4.0 |
| Precipitation (inches) | 3.2 | 3.6 | 4.0 | 3.6 | 2.8 | 6.4 | 4.2 |
| Departure from Normal | 0.4 | 0.0 | 0.0 | -0.7 | -1.8 | 2.9 | 0.2 |
| Growing Degree Days (base 32) | 336 | 769 | 928 | 1112 | 1119 | 747 | 519 |
| Departure from Normal | 4 | 64 | -25 | -4 | 63 | -41 | 73 |
| Shelburne* | April | May | June | July | August | Sept | Oct |
| Average Temperature (F) | 46.1 | 61.6 | 67.8 | 73.0 | 72.0 | 61.9 | 52.9 |
| Departure from Normal | 1.3 | 5.2 | 2.0 | 2.4 | 3.2 | 1.4 | 4.8 |
| Precipitation (inches) | 2.8 | 4.4 | 3.2 | 3.8 | 2.9 | 5.36 | 5.04 |
| Departure from Normal | 0.0 | 0.9 | -0.5 | -0.4 | -1.0 | 1.72 | 1.44 |
| Growing Degree Days (base 32) | 435 | 917 | 1072 | 1271 | 1241 | 925 | 627 |
| Departure from Normal | 51 | 161 | 58 | 73 | 102 | 68 | 126 |

*Data compiled from Northeast Regional Climate Center data from weather stations in Newport, VT and Burlington, VT. Historical averages for 30 years of NOAA data (1981-2010).

At Butterworks Farm, there was no statistical difference in yield among the nutrient dense sprays for first, second or third cut hay (Tables 5-7). However, second cut yields of the treatments 'All' and PhotoMag were higher than first cut (Figure 1), whereas yields of the other treatments decreased with each cut, as more typical in Vermont. First cut crude protein of the 'All' treatment was 18.1% compared to the control of 15.1% (Table 5). Crude protein generally increased with each cut (Figure 2), except for the 'All' and PhotoMag treatments, where first cut CP was slightly higher than second cut. 'All', PhotoMag and the Phosphorus treatments had the most favorable forage quality characteristics including the lowest fiber, highest starch, total digestible nutrients, net energy for lactation and relative feed value. Starch levels for second cut hay were highest for PhotoMag and Rejuvenate treatments (Table 6). There were no significant differences between the treatments for third cut (Table 7).

Table 5. First cut hay yield and quality, Westfield, VT, 31-May 2012.

| Treatment | Height in. | DM Yield lbs. acre ⁻¹ | DM % | CP % | ADF % | NDF % | Starch % | TDN % | NEL Mcal/lb | RFV % |
|------------|---------------|-------------------------------------|---------|--------------|--------------|--------------|-------------|--------------|----------------|-------------|
| All | 20.7* | 2234 | 17.9 | 18.1* | 29.1* | 49.2* | 3.4* | 64.5* | 0.643* | 126* |
| Control | 20.2* | 2828 | 18.6 | 15.1 | 33.3 | 56.2 | 2.9 | 62.8 | 0.595 | 105 |
| MicroPak | 19.4* | 2547 | 18.5 | 17.3* | 31.1 | 52.7 | 3.1* | 63.3 | 0.618 | 114 |
| Phosphorus | 19.3* | 2520 | 17.7 | 17.1* | 29.5* | 49.7* | 3.4* | 65.3* | 0.648* | 124* |
| PhotoMag | 17.3 | 2140 | 17.7 | 18.1* | 29.8* | 48.7* | 3.7* | 64.5* | 0.645* | 126* |
| Potassium | 18.3 | 2433 | 18.1 | 16.2 | 32.0 | 53.0 | 2.7 | 62.8 | 0.613 | 113 |
| Rejuvenate | 20.6* | 2523 | 17.4 | 17.1* | 30.5* | 52.4 | 3.1 | 64.3* | 0.628* | 116* |
| Trial Mean | 19.4 | 2461 | 18.0 | 17.0 | 30.7 | 51.7 | 3.2 | 63.9 | 0.627 | 118 |
| LSD | 1.8302 | NS | NS | 1.7923 | 1.7188 | 3.4669 | 0.5812 | 1.122 | 0.0219 | 10.46 |

*Varieties with an asterisk indicate that it was not significantly different than the top performer in column (in **bold**).

NS - None of the varieties were significantly different from one another.

Table 6. Second cut hay yield and quality, Westfield, VT, 9-Jul 2012.

| Treatment | DM Yield lbs. acre ⁻¹ | DM % | CP % | ADF % | NDF % | dNDF % | Starch % | TDN % | NEL Mcal/lb | NSC % |
|------------|-------------------------------------|---------|---------|----------|----------|-----------|--------------|----------|----------------|--------------|
| All | 2288 | 25.3 | 17.8 | 29.3 | 42.8 | 27.8 | 3.35 | 64.3 | 0.665 | 15.7 |
| Control | 2191 | 26.2 | 17.7 | 28.3 | 42.9 | 28.8 | 3.43 | 65.1 | 0.675 | 15.6 |
| MicroPak | 2255 | 25.8 | 18.3 | 28.4 | 42.3 | 26.8 | 3.45 | 65.1 | 0.675 | 16.2* |
| Phosphorus | 2177 | 25.0 | 17.5 | 29.2 | 43.9 | 28.2 | 3.38 | 64.4 | 0.668 | 15.7 |
| PhotoMag | 2391 | 25.5 | 18.1 | 27.9 | 40.8 | 25.6 | 3.83* | 65.2 | 0.675 | 16.7* |
| Potassium | 2215 | 26.4 | 17.0 | 29.3 | 44.2 | 27.8 | 3.40 | 64.8 | 0.670 | 15.9 |
| Rejuvenate | 1996 | 24.4 | 18.6 | 27.6 | 40.2 | 25.3 | 3.63* | 65.4 | 0.678 | 16.9* |
| Trial Mean | 2216 | 25.5 | 17.8 | 28.5 | 42.4 | 27.2 | 3.49 | 64.9 | 0.672 | 16.1 |
| LSD | NS | NS | NS | NS | NS | NS | 0.2529 | NS | NS | 0.8494 |

*Varieties with an asterisk indicate that it was not significantly different than the top performer in column (in **bold**).

NS - None of the varieties were significantly different from one another.

Table 7. Third cut hay yield and quality, Westfield, VT, 21-Aug 2012.

| Treatment | DM Yield lbs. acre ⁻¹ | DM % | CP % | ADF % | NDF % | dNDF % | Starch % | TDN % | NEL Mcal/lb | NSC % |
|------------|-------------------------------------|---------|---------|----------|----------|-----------|-------------|----------|----------------|----------|
| All | 1525 | 19.6 | 20.1 | 29.6 | 42.0 | 62.5 | 3.3 | 63.2 | 0.653 | 13.6 |
| Control | 1453 | 18.8 | 21.1 | 28.3 | 39.4 | 65.6 | 3.6 | 63.5 | 0.655 | 14.0 |
| MicroPak | 1166 | 17.3 | 21.2 | 28.4 | 40.2 | 64.8 | 3.3 | 63.6 | 0.658 | 13.7 |
| Phosphorus | 1297 | 19.6 | 19.8 | 29.3 | 42.2 | 64.6 | 3.2 | 63.4 | 0.658 | 13.6 |
| PhotoMag | 1477 | 18.6 | 20.8 | 29.0 | 40.6 | 64.4 | 3.5 | 63.3 | 0.653 | 13.7 |
| Potassium | 1253 | 19.3 | 20.6 | 29.3 | 41.4 | 63.8 | 3.5 | 63.0 | 0.650 | 13.5 |
| Rejuvenate | 1614 | 21.7 | 19.9 | 28.8 | 41.5 | 63.7 | 3.4 | 63.7 | 0.660 | 14.1 |
| Trial Mean | 1398 | 19.3 | 20.5 | 28.9 | 41.0 | 64.2 | 3.4 | 63.4 | 0.655 | 13.7 |
| LSD | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS |

*Varieties with an asterisk indicate that it was not significantly different than the top performer in column (in **bold**).

NS - None of the varieties were significantly different from one another.




Figure 1. First, second and third cut dry matter yields, Westfield, VT.




Figure 2. First, second and third cut crude protein, Westfield, VT, 2012. Treatments with the same letter did not differ significantly from one another.

At Shelburne Farms, there were no significant differences for yield or quality between the nutrient dense spray treatments for first, second or third cut hay (Tables 8-10). The only exception to this was second cut starch levels; the Control and Phosphorus treatments had higher starch levels than the other treatments (Table 9). Dry matter yields were very high for first cut, averaging 3097 lbs acre⁻¹, and they decreased with each subsequent cut (Figure 3). Crude protein levels were also high for first cut, averaging 17.5% (Figure 4). High yield and protein levels may be attributed to the early harvest on 17-May. Above average temperatures in March, April and May provided good conditions for plant growth and soil drying to allow for the early harvest.

Table 8. First cut hay yield and quality, Shelburne, VT, 17-May 2012.

| Treatment | Height in. | DM Yield lbs. acre ⁻¹ | DM % | CP % | ADF % | NDF % | Starch % | TDN % | NEL Mcal/lb |
|------------|---------------|-------------------------------------|---------|---------|----------|----------|-------------|----------|----------------|
| All | 27.9 | 2977 | 18.0 | 17.6 | 31.8 | 57.7 | 2.0 | 63.0 | 0.588 |
| Control | 33.1 | 3137 | 18.4 | 17.4 | 30.7 | 56.3 | 1.2 | 62.3 | 0.588 |
| MicroPak | 28.2 | 3193 | 17.8 | 16.3 | 32.0 | 57.6 | 1.6 | 62.3 | 0.583 |
| Phosphorus | 31.2 | 3030 | 17.0 | 17.4 | 30.8 | 55.4 | 1.3 | 62.3 | 0.595 |
| PhotoMag | 33.5 | 3377 | 16.5 | 17.9 | 32.2 | 58.1 | 1.2 | 61.3 | 0.568 |
| Potassium | 27.6 | 3057 | 17.5 | 17.7 | 31.4 | 55.9 | 1.9 | 62.5 | 0.590 |
| Rejuvenate | 32.8 | 2906 | 17.9 | 18.3 | 31.2 | 58.1 | 1.8 | 62.0 | 0.575 |
| Trial Mean | 30.6 | 3097 | 17.6 | 17.5 | 31.4 | 57.0 | 1.6 | 62.2 | 0.584 |
| LSD | NS | NS | NS | NS | NS | NS | NS | NS | NS |

NS - None of the varieties were significantly different from one another.

Table 9. Second cut hay yield and quality, Shelburne, VT, 21-Jun 2012.

| Treatment | Height in. | DM Yield lbs. acre ⁻¹ | DM % | CP % | ADF % | NDF % | dNDF % | Starch % | TDN % | NEL Mcal/lb | NSC % |
|------------|---------------|-------------------------------------|---------|---------|----------|----------|-----------|-------------|----------|----------------|----------|
| All | 17.2 | 2369 | 27.8 | 14.9 | 35.9 | 55.4 | 63.8 | 3.0 | 59.8 | 0.615 | 10.0 |
| Control | 15.8 | 2379 | 25.2 | 15.1 | 35.3 | 53.8 | 65.2 | 3.3* | 59.9 | 0.618 | 8.4 |
| MicroPak | 17.6 | 2209 | 26.0 | 14.8 | 36.0 | 56.1 | 64.8 | 3.0 | 59.7 | 0.613 | 9.7 |
| Phosphorus | 16.9 | 2253 | 27.0 | 14.6 | 35.8 | 56.0 | 64.8 | 3.1* | 59.7 | 0.613 | 9.8 |
| PhotoMag | 17.3 | 2663 | 26.5 | 14.9 | 36.4 | 55.7 | 63.0 | 2.9 | 59.0 | 0.605 | 9.5 |
| Potassium | 17.6 | 2322 | 27.7 | 14.2 | 36.2 | 56.7 | 64.9 | 3.0 | 59.6 | 0.613 | 9.8 |
| Rejuvenate | 16.3 | 2309 | 28.1 | 14.8 | 36.2 | 55.8 | 64.7 | 2.9 | 59.3 | 0.608 | 9.4 |
| Trial Mean | 16.9 | 2358 | 26.9 | 14.7 | 36.0 | 55.6 | 64.5 | 3.0 | 59.5 | 0.612 | 9.5 |
| LSD | NS | NS | NS | NS | NS | NS | NS | 0.1978 | NS | NS | NS |

*Varieties with an asterisk indicate that it was not significantly different than the top performer in column (in **bold**).

NS - None of the varieties were significantly different from one another.

Table 10. Third cut hay yield and quality, Shelburne, VT, 27-Jul 2012.

| Treatment | DM Yield lbs. acre ⁻¹ | DM % | CP % | ADF % | NDF % | dNDF % | Starch % | TDN % | NEL Mcal/lb | NSC % |
|------------|-------------------------------------|---------|---------|----------|----------|-----------|-------------|----------|----------------|----------|
| All | 1366.3 | 27.7 | 18.6 | 28.7 | 48.2 | 59.2 | 2.4 | 62.8 | 0.650 | 9.8 |
| Control | 1246.4 | 26.7 | 18.9 | 28.0 | 46.9 | 58.9 | 2.5 | 63.1 | 0.653 | 9.9 |
| MicroPak | 1365.6 | 27.1 | 18.2 | 29.2 | 49.1 | 60.3 | 2.3 | 62.5 | 0.650 | 9.5 |
| Phosphorus | 1297.4 | 27.0 | 18.1 | 29.2 | 49.5 | 60.1 | 2.3 | 62.3 | 0.645 | 9.8 |
| PhotoMag | 1518.5 | 25.9 | 19.1 | 28.4 | 47.8 | 60.4 | 2.3 | 62.6 | 0.648 | 9.5 |
| Potassium | 1300.6 | 25.9 | 18.7 | 29.2 | 48.5 | 59.5 | 2.0 | 62.3 | 0.645 | 9.6 |
| Rejuvenate | 1286.2 | 27.3 | 18.6 | 28.4 | 48.3 | 59.7 | 2.3 | 63.0 | 0.650 | 9.4 |
| Trial Mean | 1340.1 | 26.8 | 18.6 | 28.7 | 48.3 | 59.7 | 2.3 | 62.7 | 0.649 | 9.6 |
| LSD | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS |

NS - None of the varieties were significantly different from one another.




Figure 3. First, second and third cut dry matter yields, Shelburne, VT, 2012.




Figure 4. First, second and third cut crude protein, Shelburne, VT, 2012.

In addition, there was no significant difference among the treatments in calcium, phosphorus, magnesium, potassium or sulfur at either location (data not shown).

ACKNOWLEDGEMENTS

The UVM Extension Northwest Crops and Soils team would like to thank Butterworks Farm and Shelburne Farms for hosting these trials, and the Lattner Foundation for funding this research.

UVM Extension helps individuals and communities put research-based knowledge to work.

Any reference to commercial products, trade names, or brand names is for information only, and no endorsement or approval is intended. Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. University of Vermont Extension, and U.S. Department of Agriculture, cooperating, offer education and employment to everyone without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or familial status.

