

University of Vermont

UVM ScholarWorks

Family Medicine Clerkship Student Projects

Family Medicine Community

2019

Intimate Partner Violence: Improving Screening Rates in the Primary Care Setting

Alexandra E. Churchill
University of Vermont

Follow this and additional works at: <https://scholarworks.uvm.edu/fmclerk>




Part of the [Medical Education Commons](#), and the [Primary Care Commons](#)

Recommended Citation

Churchill, Alexandra E., "Intimate Partner Violence: Improving Screening Rates in the Primary Care Setting" (2019). *Family Medicine Clerkship Student Projects*. 475.
<https://scholarworks.uvm.edu/fmclerk/475>

This Book is brought to you for free and open access by the Family Medicine Community at UVM ScholarWorks. It has been accepted for inclusion in Family Medicine Clerkship Student Projects by an authorized administrator of UVM ScholarWorks. For more information, please contact scholarworks@uvm.edu.



INTIMATE PARTNER VIOLENCE: IMPROVING SCREENING RATES IN THE PRIMARY CARE SETTING

Newtown Primary Care
Lexi Churchill
Rotation 1 - April 2019
Dr. Eurica Chang

PROBLEM IDENTIFICATION AND DESCRIPTION OF NEED

Intimate partner violence (IPV) is defined by the CDC as “physical violence, sexual violence, stalking and psychological aggression (including coercive acts) by a current or former intimate partner”¹

- ❖ National data (2015) indicate 1 in 4 women and 1 in 10 men have experienced contact sexual violence, physical violence, and/or stalking with subsequent IPV-related impact²
 - ❖ Greater than one-third of both women and men have experienced psychological aggression alone²
 - ❖ 71.1% of female victims and 55.8% of male victims first experienced IPV before age 25²
- ❖ Domestic violence agencies across Connecticut provided services to 38,192 victims in 2018³
- ❖ Though the U.S. Preventative Services Task Force recommends screening all female patients of childbearing age for IPV³, primary care provider implementation rates are low⁴
- ❖ Within the Western Connecticut Health Network EMR, none of the publicly available “Wellness Visit” templates include IPV screening, nor is there a separate IPV screening template

PUBLIC HEALTH COST

- ❖ Outside of direct physical injury, IPV is associated with numerous medical conditions including cardiovascular disease, CNS and GI disorders, chronic pain syndromes, anxiety, depression, and PTSD¹
- ❖ Victims of IPV are at increased risk of miscarriage, preterm labor, STIs, high-risk sexual behavior, homelessness, substance abuse, poor nutrition, alcoholism, and suicide¹
- ❖ One study estimated the lifetime economic burden of IPV victims, both male and female, to be nearly \$3.6 trillion, of which \$2.1 trillion (59%) came from medical costs⁵
- ❖ Under-reporting of IPV by victims indicates the actual costs are likely even higher

COMMUNITY PERSPECTIVE

Nya Rossi, PA *Newtown Primary Care*

What are the barriers facing patients to disclose IPV to their primary care provider?

Fear. Of my patients that are known victims of IPV, their partners usually come to the office visits, making it difficult to directly ask the patient about IPV or to offer resources discretely.

What other barriers keep providers from asking screening IPV questions?

It is uncomfortable to ask. I was trained to ask 'Do you feel safe in your relationship?', but that doesn't seem adequate and wording can be difficult with such a sensitive topic. I should be asking it more often though. Including a well-worded, one-question screen in annual visits would be a great way to ensure the question gets asked.

LCSW *Western Connecticut Health Network*

What is your perspective of IPV screening during routine office visits?

It's absolutely necessary, it helps normalize the question and provides an avenue for victims to safely disclose without fear of repercussions. Though it can be a tough question to ask because it seems so direct, so it's probably not happening enough. I will see patients for one mental health issue – depression, anxiety, PTSD, etc. – and it will come out at a later session that the patient is a victim of IPV. I don't think their provider knows at the time they give the referral for them to come talk to me.

How do you think rates of provider screening for IPV might be improved?

Just by making it a habit to follow up the routine questions about sexual health with asking about the patient's relationships – how they're doing, then naturally flows into if the patient has ever felt unsafe.

INTERVENTION AND METHODOLOGY

- ❖ Newtown Primary Care providers most frequently utilized HPI templates adapted from pre-existing public phrases within the EMR system to structure interviews during wellness visits
- ❖ **Intervention:** Create a concise, effective IPV screening template for use within annual wellness visit HPI phrases to increase provider awareness, ease of access, and implementation
- ❖ USPSTF recommends initial screening with a validated tool, followed by referral to intervention services, including local and national resources, if the screen is positive⁴
 - ❖ Evaluation of the validated screening tools identified the Woman Abuse Screening Tool – Short Form (WAST-SF) as time efficient and efficacious (sensitivity: 93%, specificity: 68%)⁷
- ❖ Positive screens should be assessed for risk of immediate harm⁶

RESULTS

❖ IPV screening template (right) comprised of the WAST-SF, a 5-question immediate harm risk follow up, and resources was disseminated among the Newtown Primary Care providers

❖ 3 of the 6 providers self-reported incorporating the IPV screening tool into their personalized HPI templates for annual wellness visits

IPV Screening Template:

Are you currently in a relationship? **No/Yes**

In general, how would you describe your relationship? **No tension/Some tension/A lot of tension**

You and your partner work out arguments with: **No difficulty/Some difficulty/Great difficulty**

If patient responded **A lot of tension** or **Great difficulty**, assess immediate harm risk:

Has the physical violence increased over the past six months? **No/Yes**

Has your partner used a weapon or threatened you with a weapon? **No/Yes**

Do you believe your partner is capable of killing you? **No/Yes**

Have you been beaten while pregnant? **No/Yes**

Is your partner violently and constantly jealous of you? **No/Yes**

Resources were **not applicable**/provided/deferred, including follow up with in-house Behavioral Health services, Women's Center of Greater Danbury and National Domestic Violence Hotline 1-800-799-SAFE (7233).

Blue text indicates default selections from an EMR drop-down menu

EVALUATION OF EFFECTIVENESS AND LIMITATIONS

- ❖ IPV screening template was met with positive feedback from the Newtown Primary Care office providers
- ❖ Inclusion of the screen into providers' HPI templates increases provider awareness at a minimum, even if the questions are not asked
 - ❖ Follow-up quantitative data needed to assess provider screening rates
- ❖ Consistent use of the amended HPI template facilitates provider comfort with screening and normalizes screening questions at wellness visits
- ❖ Inability to create a network-wide public phrase severely minimized ubiquity of the screening tool in WCHN
 - ❖ Currently template is only available to providers it has been personally shared with
- ❖ Difficulty implementing the screen if partner is present during the wellness visit
- ❖ Time constraints limited evaluation of screening implementation and patient response

RECOMMENDATIONS FOR FUTURE INTERVENTIONS

- ❖ Work with EMR technology services to incorporate the IPV screening tool directly into preexisting public phrases or create an independent IPV screening public phrase for network-wide provider access
- ❖ Regularly assess provider screening rates through QA initiatives
- ❖ Coordinate with the Women's Center of Greater Danbury to organize staff training to reinforce provider awareness and provide tools for a conscientious approach to IPV screening
 - ❖ Incorporate a one-on-one portion of wellness visits in which to ask sensitive screening questions
- ❖ Place IPV information in the office lobby, restrooms, and exam rooms to facilitate community awareness and discrete dissemination of resources
- ❖ Gauge patient response to IPV screening via surveys

REFERENCES

1. *Preventing Intimate Partner Violence*. Centers for Disease Control and Prevention. 2019. 4/22/2019. <https://www.cdc.gov/violenceprevention/intimatepartnerviolence/definitions.html>
2. Smith, S.G., Zhang, X., Basile, K.C., Merrick, M.T., Wang, J., Kresnow, M., Chen, J. (2018). *The National Intimate Partner and Sexual Violence Survey (NISVS): 2015 Data Brief – Updated Release*. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
3. *Connecticut Service Statistics*. Connecticut Coalition Against Domestic Violence. 2018. 4/22/2019. <http://www.ctcadv.org/information-about-domestic-violence/statistics/>
4. Moyer VA. *Screening for intimate partner violence and abuse of elderly and vulnerable adults: U.S. Preventive Services Task Force recommendation statement*. *Ann Intern Med*. 2013;158(6):478–486.
5. Peterson C, Kearns MC, McIntosh WL, Estefan LF, Nicolaidis C, McCollister KE, Gordon A, Florence C. Lifetime Economic Burden of Intimate Partner Violence Among U.S. Adults. *Am J Prev Med*. 2018 Oct;55(4):433-444. doi: 10.1016/j.amepre.2018.04.049. Epub 2018 Aug 22. PubMed PMID: 30166082; PubMed Central PMCID: PMC6161830.
6. Dicola D, Spaar E. *Intimate partner violence*. *Am Fam Physician*. 2016;94(8):646-651.
7. Rabin R, Jennings J, Campbell J, Blair-Merritt M. Intimate partner violence screening tools. *Am J Prev Med*. 2009;36(5):439- 445