University of Vermont UVM ScholarWorks

Graduate College Dissertations and Theses

Dissertations and Theses

2021

Unique Genomic And Phenotypic Responses To Extreme And Variable Ph Conditions In Purple Urchin Larvae, Strongylocentrotus Purpuratus

April Dawn Garrett University of Vermont

Follow this and additional works at: https://scholarworks.uvm.edu/graddis

Part of the Biology Commons

Recommended Citation

Garrett, April Dawn, "Unique Genomic And Phenotypic Responses To Extreme And Variable Ph Conditions In Purple Urchin Larvae, Strongylocentrotus Purpuratus" (2021). *Graduate College Dissertations and Theses*. 1335.

https://scholarworks.uvm.edu/graddis/1335

This Thesis is brought to you for free and open access by the Dissertations and Theses at UVM ScholarWorks. It has been accepted for inclusion in Graduate College Dissertations and Theses by an authorized administrator of UVM ScholarWorks. For more information, please contact scholarworks@uvm.edu.

UNIQUE GENOMIC AND PHENOTYPIC RESPONSES TO EXTREME AND VARIABLE PH CONDITIONS IN PURPLE URCHIN LARVAE, STRONGYLOCENTROTUS PURPURATUS

A Thesis Presented

by

April D. Garrett

to

The Faculty of the Graduate College

of

The University of Vermont

In Partial Fulfillment of the Requirements for the Degree of Master of Science Specializing in Biology

January, 2021

Defense Date: November 12, 2020 Thesis Examination Committee:

Melissa H. Pespeni, Ph.D., Advisor Stephen R. Keller, Ph.D., Chairperson Brent L. Lockwood, Ph.D. Alicia M. Ebert, Ph.D. Cynthia J. Forehand, Ph.D., Dean of the Graduate College

ABSTRACT

Given the unprecedented increases in atmospheric carbon dioxide and its projected negative impacts on organismal ecology and physiology, it is crucial to understand if and how organisms will withstand such environmental changes. Due to the oceans' service as a carbon sink, marine organisms face the added stressor of ocean acidification (OA), the process by which carbon dioxide mixes with water and decreases pH while simultaneously depleting the seawater of calcium carbonate. Marine organisms that rely on calcium carbonate for exoskeleton development are considered particularly vulnerable to OA, though previous results vary among species, leading to the question of who the real 'winners' and 'losers' will be in the face of increasing OA. Strongylocentrotus *purpuratus*, the purple sea urchin, is one such calcifying organism whose ability to respond to OA is relatively well studied in the past decade, but its future success still remains largely unclear. Within their natural habitat of the California Current Marine Ecosystem (CCME), there exists not only more extreme mean sea surface pH values as compared to the open ocean, but also high spatial and temporal variability due to a natural phenomenon known as upwelling. My thesis research aims to use theoretical and experimental tools from population genetics, experimental evolution, and ecological genomics to determine if developing purple sea urchins have the genetic capacity and physiological capability to respond to future OA in both static and variable extreme pH conditions. Low (pH 7.5) and extreme (pH 7.0) pH conditions led to decreased survival, with variability helping recover survival in those treatments. However, this recovery came with a trade-off: survivors in the extreme variable treatment were significantly smaller in body size compared to their static counterpart. Further, my work shows that purple urchins have the genomic capacity to respond uniquely to both extreme and variable pH conditions. While these results may be promising for the early life stages of the purple sea urchin, the carry-over effects of future low pH in the CCME on surviving larvae undergoing metamorphosis and developing into reproductive adulthood remain to be studied, as do the responses of marine species with lower levels of standing genetic variation in the face of increasing OA and pH variability in the CCME.

CITATIONS

Material from this thesis has been published in the following form:

Garrett, A. D.; Brennan, R.S.; Steinhart, A.L.; Pelletier, A.M.; Pespeni, M.H.. (2020). Unique genomic and phenotypic responses to extreme and variable pH conditions in purple sea urchin larvae. Journal of Integrative and Comparative Biology, 60, 318-331.

DEDICATIONS

I dedicate this thesis to the women in my life who have made me the woman I am today spiritually, personally, and/or academically: my dear and most wonderful mother Jamie S. Mayer ('Momma'), my beloved and missed grandmothers, Eloise M. Terry and Dorothy 'Dottie' J. Fay, my 'spiritual mom' and dear friend Kerri A. Jones, and my most amazing graduate and undergraduate advisors who have inspired me beyond words and will continue to do so every day of my life, Melissa H. Pespeni and Cheryl A. Logan. Thank you all, from the bottom of my heart, for being my role models and for your endless encouragement and support.

ACKNOWLEDGEMENTS

I am extremely grateful for all of the love, encouragement, and support shown to me over the years and would like to take this moment to thank several people. First and foremost, I thank God, for I know I would not be here without Him and am so grateful for the life and opportunities I have had both within and outside of academia. The good times have brought me smiles and laughter, and the bad times have helped shape and refine me into who I am today as both a young woman and a scientist. I also thank my 'Momma' for loving me unconditionally and always supporting and praying for me. You are my role model, and I love you dearly. You are without a doubt the best mother ever. Thank you to all of my parents for believing in me and encouraging me to do and be my best. To Joseph and Kayla: thank you for your love and support. You both have believed in me and helped motivate me to successfully complete this thesis, along with our parents, and I'm grateful for you both and love all of my family very much. Then there are the friends who I chose to be family and who chose me to be theirs. Kevin, you are one of my best friends and have changed my life for the better; I cherish and respect you greatly. Thank you for your grace, love, kindness, and friendship, and for bringing me to a place where I could heal and grow into the person I've always wanted to be, the person the Lord made me to be. And thank you to Sandi for your part in my healing and growth as well and for being such a great role model. Kerri, I am so blessed by your mentorship and friendship and love you dearly - thank you so much for your prayers, wisdom, kindness, and role in my growth. You have blessed my life immensely. To my other best friends – Sarah, Kami, Kyndra, Luiza, and Erin – I love you all like sisters and feel so blessed to know your beautiful souls. Thank you for your support and encouragement. A shoutout to the Mersino family as well - much love to 'Scobbit' and Elisabeth and your daughters. Thank you for your sweet friendships. Thank you to all of my friends who have invested in my life and allowed me to invest in theirs. I thank my church family as well, both the SBC Open Table and Q.U.E.E.N.S. groups for your prayers, encouragement, and support both in my thesis as well as in my personal life. You are like family; thank you for blessing my life (and a special

shoutout to Kelsey). Thank you to my East Coast family (Leigh, Ellen, and Laura) and my East Coast grandparents (Bill & Jean). You've all loved and supported me throughout the years in VT as well as after I left, and I love you all very much and look forward to my next visit. To my other dear VT friends, Rebecca and Caitlin & Chris - I love you all, I cherish our memories together, and I look forward to a future VT visit again someday. Thank you to my colleagues in the Pespeni lab and at my current workplace for your support over the years. You are all amazing scientists that I've had the privilege to get to work with, and I have a lot of respect for all of you. Further, I thank my coauthors for their contributions in publishing this thesis research. Dr. Reid Brennan helped with coding, data analysis, and manuscript revisions, and was always available to answer questions and discuss ideas. Anya Steinhart and Aubrey Pelletier helped with experimental implementation and collection and analysis for the morphometric and survival data, respectively. Finally, I thank my wonderful graduate advisor, Dr. Melissa Pespeni, who helped with experimental design and implementation, data analysis, and manuscript revisions. I remember my first day in your office as your first graduate student – getting to work with you was so unreal to me. I remember telling you I felt like I was talking to a rockstar. Here we are all these years later, and I still see you this way. You are an inspiring and absolutely amazing scientist. Thank you for everything you've done for me – I will never forget.

TABLE OF CONTENTS

Pag	e
DEDICATIONSiii	
ACKNOWLEDGMENTSiv	
INTRODUCTORY LITERATURE REVIEW1	
CHAPTER 1: UNIQUE GENOMIC AND PHENOTYPIC RESPONSES TO	
EXTREME AND VARIABLE PH CONDITIONS IN PURPLE SEA URCHIN	
LARVAE	
BIBLIOGRAPHY	,

INTRODUCTORY LITERATURE REVIEW

Increases in atmospheric carbon dioxide since pre-industrial times has consequently led to global changes in our climate, including more frequent climate extremes as well as increases in variability (IPCC 2020, Rackley 2017, Vázquez et al., 2017), the latter which is thought to be of even more importance than simply increased averages of climate variables (Katz & Brown 1992). Such drastic changes in an organisms' environment can lead to physiological and evolutionary consequences (Vázquez et al., 2017). On the other hand, experiencing more extreme and variable environmental conditions can lead to genetic or physiological priming of individuals and subsequent growth or persistence of their populations, serving as a form of evolutionary rescue (Carlson et al., 2014) and leading to rapid adaptation if enough natural standing genetic variation exists in a population (Lande & Shannon 1996). In order for this rapid adaptation to occur and ultimately for populations to persist, early life history stages of the organisms that make up the population must be able to reach sexual maturity and reproduce, a situation that is further complicated when organisms are placed under more extreme and varying environmental conditions, particularly those outside of their natural range. This complication is due to the general phenotypic link between early and later life history stages, or 'carry-over' effects, where the phenotype of earlier developmental stages has an impact on the later life history stages of an organism, such as seen in marine invertebrates that experience different and more extreme dynamics as larvae in the pelagic before settling down as relatively sessile adults in the benthic (Marshall & Morgan 2011). Any negative impacts from extreme or variable environmental events on

earlier life history stages can have downstream negative consequences on later developmental stages, leading to a decline in reproductive individuals within a population and ultimately threatening population persistence.

Given the unprecedented increases in atmospheric carbon dioxide and its projected negative impacts on organismal ecology and physiology, it is crucial to understand if and how organisms of different life history stages will withstand such environmental changes. Ocean acidification (OA) is one such resultant global change phenomenon that marine invertebrates of different life history stages experience, where carbon dioxide mixes with sea surface water and decreases its pH while simultaneously depleting the seawater of natural calcium carbonate (Doney et al., 2009). The associated decrease in pH has been shown to disrupt acid-base physiology in a variety of marine organisms (Fabry et al., 2008, Esbaugh et al., 2012), and the simultaneous depletion of calcium carbonate exacerbates the negative impacts of OA particularly on calcifying marine invertebrates that rely on naturally available carbonate to grow and maintain structural integrity. This has been shown to be particularly difficult for early life stages of calcifying marine invertebrates (Kurihara 2008, Kroeker et al., 2010). Similar to the concept of trying to determine the 'winners' and 'losers' under increasing climate change (Somero 2010), it is important to determine which calcifying species, if any, will be able to 'win' against an ocean increasing in its absorption of carbon dioxide. While migration and acclimatization are two ways in which some species may be able to tolerate current increases in OA, the real 'winners' under increasing OA will be those who can survive early life history stages and adapt at a rapid enough pace, as the oceans are not only

increasing in acidity over time, but also the rate of increase of OA itself is increasing (Doney et al., 2009).

Rapid adaptation in an acidifying ocean, as with any other selective pressure, requires the existence of ample standing genetic variation underlying phenotypic variation on which selection can act (Sunday et al., 2014). In order for evolution to occur, this variation needs to be heritable (Lynch & Walsh 1998, Sunday et al., 2014) and will result in differential fecundity, whereby some organisms within a population will have alleles that allow their survival in lower pH and/or carbonate saturation and those individuals will go on to reproduce and pass on these alleles to the next generation. Thus, the frequency of these alleles in the population will change over time, a standard definition of evolution in population genetic theory (Nielsen & Slatkin 2013), with a presumed increase in frequency of alleles that are adaptive. Relatively few studies have measured changes in allele frequencies in response to selective pressures in natural populations to link adaptive phenotypes to genotypes (Schlötterer et al., 2015). For example, few but significant changes in allele frequencies were found in stick insects transplanted to different host plants, allowing for particular genomic loci under selection to be detected, in addition to observed phenotypic changes in color (Gompert et al., 2014). Genomic targets of selection have also been determined and functionally validated in Drosophila that underlie their adaptation to Drosophila C virus (Martins et al., 2014). Further, recent studies have shown rapid selection in action linked to underlying genomic loci, with increases in allele frequency of a mutation linked to survival in mice across 14 months (Barrett et al., 2019) and genomic differentiation and signatures of local

adaptation in several loci involved in nervous system functions in green anole lizards after just one extreme winter season (Campbell-Staton et al., 2017).

In marine organisms, few studies to date have measured allele frequency changes or identified genetic targets of selection in response to OA. However, of the work that has been done, significant allele frequency changes have been found for loci in genes involved in oxidative phosphorylation, RNA transcription, and ribosomal structure in copepods and biomineralization, ion homeostasis, and lipid metabolism in urchins (De Wit et al., 2016, Pespeni et al., 2013a, Brennan et al., 2019). For most marine organisms experiencing OA as a selective pressure, their ability to rapidly adapt is challenged by their longer generation times, where the next reproductive generation may not come for 2-3 years at least. With high standing genetic variation, though, and high levels of gene flow coupled with selection, adaptive alleles for OA can be maintained in natural populations (Pespeni et al., 2013b, Pespeni and Palumbi 2013), though concern remains as to whether or not marine organisms can adapt fast enough to keep pace with increasing global change (Calosi et al., 2016). Thus, studying a marine invertebrate that is likely to be negatively impacted by OA (i.e., marine calcifiers) while also having a long generation time makes it highly challenging to conduct evolution experiments in the lab, at least in terms of looking for genetic variation that is heritable or changes in allele frequencies over multiple generations (Sunday et al., 2014). However, while accurately being able to estimate heritability of traits and additive genetic variance is rather infeasible with most marine life (Lynch & Walsh 1998, Kelly et al., 2013), previous work has shown that experimental selection can be conducted in the lab for a marine calcifier, the purple sea urchin (Strongylocentrotus purpuratus) within a single generation,

allowing for the identification of genes targeted by selection in response to acidification (Pespeni et al., 2013a, Brennan et al., 2019) and thus a better understanding of how some marine species may adapt to increasing OA.

The purple sea urchin provides a great system for determining targets of selection at early life history stages in a marine calcifier responding to extreme and varying OA, having served for many years as a model 'non-model' marine species. S. *purpuratus* is a key grazer and kelp forest ecosystem engineer as well as a valuable fishery (Pearse 2006). In addition to its ecological and economic values, the purple sea urchin is increasingly used for OA studies due to its residence within the California Current Marine Ecosystem (CCME), renowned for its lower and more variable pH conditions from a natural phenomenon known as upwelling, and within this system has a wide geographic range extending from Alaska to Baja California, Mexico (Rogers-Bennett 2007). With such a broad range in this current system, there is a high amount of gene flow present, such that broadscale population structure is essentially nonexistent (Palumbi & Wilson 1990). One reason it has been so useful over the past couple decades in evolutionary and developmental biology studies is its abundant genetic resources that are available, with its genome published since 2006 (Sodergren et al., 2006). Additionally, the purple urchin genome is highly polymorphic (4-5% genome-wide variation, Sodergren et al., 2006) and their standing genetic variation high in nature, providing ample genomic loci on which selection can act and thus a useful system for understanding molecular mechanisms important for adaptation. Additionally, the ease of bringing urchins into a lab, spawning them to release millions of eggs and sperm, and controlling fertilization crosses allows for the capturing of their natural genetic variation

in the lab and the ability to create replicate populations for experimental evolution studies, as urchin larvae are small and allow for a large N in replicate culturing vessels. Methods have also been established for growing purple urchin larvae and inducing metamorphosis (Strathmann 2017, Pespeni et al., 2013a), allowing for the study of different critical early life history stages. This is important in terms of studying a marine calcifier's response to OA, since early life history stages of calcifiers may be more sensitive when responding to decreases in pH and carbonate as these organisms try to develop into adulthood (Kurihara 2008). All in all, this species and study system can be used to elucidate genomic and physiological capabilities of calcifying marine larvae in extreme and variable pH conditions, as experienced within the CCME and exacerbated by increasing OA.

In my thesis research, I test the hypotheses that low and extreme static pH conditions result in a decrease in pluteus larval sea urchin 1) survival and 2) growth, with pH variability rescuing the decreases seen in static pH conditions. Further, I predict that there will be both shared and unique genomic responses to static and variable pH conditions, with some loci responding uniquely to low and extreme static and variable pH conditions and not just increased or more extreme changes within the same loci.

CHAPTER 1: UNIQUE GENOMIC AND PHENOTYPIC RESPONSES TO EXTREME AND VARIABLE PH CONDITIONS IN PURPLE URCHIN LARVAE

Abstract

Environmental variation experienced by a species across space and time can promote the maintenance of genetic diversity that may be adaptive in future global change conditions. Selection experiments have shown that purple sea urchin, *Strongylocentrotus purpuratus*, populations have adaptive genetic variation for surviving pH conditions at the "edge" (pH 7.5) of conditions experienced in nature. However, little is known about whether populations have genetic variation for surviving low-pH events beyond those currently experienced in nature or how variation in pH conditions affects organismal and genetic responses. Here, we quantified survival, growth, and allele frequency shifts in experimentally selected developing purple sea urchin larvae in static and variable conditions at three pH levels: pH 8.1 (control), pH 7.5 (edge-of-range), and pH 7.0 (extreme). Variable treatments recovered body size relative to static treatments, but resulted in higher mortality, suggesting a potential tradeoff between survival and growth under pH stress. However, within each pH level, allele frequency changes were overlapping between static and variable conditions, suggesting a shared genetic basis underlying survival to mean pH regardless of variability. In contrast, genetic responses to pH 7.5 (edge) versus pH 7.0 (extreme) conditions were distinct, indicating a unique

genetic basis of survival. In addition, loci under selection were more likely to be in exonic regions than regulatory, indicating that selection targeted protein-coding variation. Loci under selection in variable pH 7.5 conditions, more similar to conditions periodically experienced in nature, performed functions related to lipid biosynthesis and metabolism, while loci under selection in static pH 7.0 conditions performed functions related to transmembrane and mitochondrial processes. While these results are promising in that purple sea urchin populations possess genetic variation for surviving extreme pH conditions not currently experienced in nature, they caution that increased acidification does not result in a linear response but elicits unique physiological stresses and survival mechanisms.

Introduction

The annual average global atmospheric carbon dioxide concentration recently reached 417ppm, likely the highest level in the past 20 million years (Rackley 2017; IPCC 2019). Consequently, global climate is changing, with alterations not only in mean conditions, but also in the variability and frequency of extreme events (Rahmstorf and Coumou 2011; Kwiatkowski and Orr 2018; McNeil and Sasse 2016). This increase in variability is driving novel conditions that exceed previous extremes (Easterling et al. 2000). For organisms to persist under these drastic changes, physiological and genetic responses will be required (Somero 2010,; Hoffmann and Sgrò 2011; Vázquez et al. 2017). However, whether populations have the adaptive potential to survive conditions beyond those currently experienced in nature remains a critical area of investigation for understanding species resilience (Lande and Shannon 1996; Flanagan et al. 2018). Further, it is unclear if environmental variation attenuates or exacerbates physiological stress and if adaptation to fluctuating versus static conditions uses the same adaptive genetic variation.

While much of our understanding of fitness in variable environments draws from studies on temperature variation (Beardmore and Levine 1963; Long 1970; Estay et al. 2011; Folguera et al. 2011; Bozinovic et al. 2011; Shama 2017), other factors are simultaneously shifting due to human activities. Oceans, in particular, are becoming more acidic due to the dissolution of atmospheric carbon dioxide into sea surface waters, declining by an approximate range of 0.017-0.027 pH units per decade since the late 1980s (IPCC 2019). This process, known as ocean acidification (OA), decreases ocean pH while simultaneously depleting the seawater of natural carbonate, the building block

for calcium carbonate (Doney et al. 2009), creating a physiologically challenging environment for many species. Maintaining acid-base balance is essential to maintain cellular functioning, and alterations to environmental pH require energetically costly intracellular compensation (Fabry et al. 2008; Esbaugh et al. 2012; Stumpp et al. 2012; Mangan et al. 2017). Organisms that develop calcareous skeletons or shells are faced with the additional difficulty of laying down and maintaining these structures under biochemically unfavorable conditions, which can lead to negative impacts on growth and survival (Byrne et al. 2013; Kroeker et al. 2010). However, some marine environments experience natural fluctuations in pH across space and time. In the California Current Marine Ecosystem (CCME), upwelling can drive diurnal fluctuations of up to 0.67 pH units, reaching the low open ocean pH levels predicted for the end-of-the-century (Yu et al. 2011; Evans et al. 2013; Chan et al. 2017; IPCC 2019). Coupled with increasing atmospheric carbon dioxide, sea surface waters within the CCME frequently experience lower pH conditions than 8.1, the open ocean average (Chan et al. 2017). Consequently, populations within this system harbor physiological mechanisms and genetic variation to tolerate low pH events (Evans et al. 2013; Pespeni et al. 2013a; Brennan et al. 2019), making them an ideal model to understand how adaptation will likely proceed as pH continues to decrease.

Our understanding of how variable pH conditions that mirror natural conditions may impact species persistence and performance is limited. Previous work has shown both negative and positive effects of pH fluctuations on organismal performance and appears to depend on factors such as the species, population origin, or trait of interest. Particularly interesting is a potential rescue effect, or mitigation, where pH variability may improve a suite of characteristics including behavior in non-calcifiers (Jarrold et al. 2017) and survivorship (Dufault et al. 2012) and growth in calcifiers (Frieder et al. 2014). Conversely, growth in some calcifying species has been shown to be negatively impacted by varying pH (Li et al. 2016, Price et al. 2012). Thus, while impacts of static low pH on marine calcifiers are generally accepted as negative, impacts of fluctuating pH appear to be less conclusive. For most calcifying species that reside specifically in pH-fluctuating environments like the CCME, questions remain as to what role naturally-varying pH will play in species' responses to OA as they reach the edge of their current natural low pH range and more rapidly and frequently encounter extreme levels beyond their current range.

We focus on the ecologically and economically important calcifying species resident to the CCME, the purple sea urchin, *Strongylocentrotus purpuratus*. This species inhabits a broad range across the west coast of North America where it experiences natural fluctuations in pH geographically and temporally (Evans et al. 2013; Pespeni et al. 2013b), with diurnal fluctuations as great a range as 0.8 pH units and lowest pH values measured at 7.43 pH units (Chan et al. 2017). Previous work has shown that *S. purpuratus* harbors adaptive standing genetic variation to rapidly adapt to low pH conditions that fall within the range typically experienced in nature (Pespeni et al. 2013a; Brennan et al. 2019). Further, this species can be readily spawned in the lab, generating hundreds of thousands of offspring harnessing the high genetic variation found in the wild. By rearing replicate pools of these larvae in a selective environment, we can identify genetic variants that shift in frequency consistently across replicates, thus identifying the standing genetic variation and related physiological functions that enable

survival across a single generation (<u>Pespeni et al. 2013a</u>; <u>Brennan et al. 2019</u>). Single generation experiments are extremely useful for this long-lived species that requires two years before reaching reproductive maturity (<u>Leahy 1986</u>).

Here, we use *S. purpuratus* to conduct a single-generation selection experiment to identify the genetic variation underlying adaptation to static and fluctuating low pH conditions. Specifically, we leverage variable and static selection regimes at pH conditions that fall within and outside the range typically experienced by this species in the wild to address the following objectives: (1) test the prediction that variability in low and extreme pH 'rescues' larval phenotypes, resulting in higher larval survival and larger body size compared to their static counterparts, (2) determine if mean pH, regardless of variability, drives selective responses, and (3) determine if larvae can use the same genomic and physiological machinery to survive as pH stress extends beyond the range experienced in nature.

Methods

Sea urchin collection & experiment

In June 2018, 25 adult *S. purpuratus* (14 females and 11 males) were collected from San Diego, CA and shipped overnight to the University of Vermont. While the frequency and magnitude of low static and fluctuating pH events in the southern part of the CCME is notably less than that of more northern regions (Hofmann et al. 2011, Yu et al. 2011; Evans et al. 2013), previous work has shown that purple urchin have high gene flow of low pH-adaptive genetic variants throughout the CCME (Palumbi and Wilson 1990). Further, recent work utilizing a similar experimental system and sourcing adults from the same location has shown there are low pH-adaptive alleles for responding to levels at the edge of this species' low pH range (Brennan et al. 2019). Immediately upon arrival, adults were induced to spawn by injecting 0.5M KCl into the peristome (Strathmann 2017, Brennan et al. 2019). Eggs were filtered over 215-micron mesh and density was determined to partition 70,000 eggs from each female into each of 3 static pH conditions: 8.1, 7.5, and 7.0 (210,000 eggs/female). The first two pH conditions were chosen to represent different degrees of pH in the environment frequently encountered by this species in the wild. The control pH condition (8.1) represents a benign treatment that reflects the current average open ocean pH conditions (Chan et al. 2017). Similarly, the intermediate pH condition (7.5) is at the edge of the pH range that purple sea urchin naturally experience in the CCME (Evans et al. 2013; Chan et al. 2017), but is also equal to the predicted end-of-century open ocean pH levels should carbon emissions continue 'business as usual' (IPCC 2019). The extreme low pH condition (7.0) falls outside of the range typically experienced in the wild. However, continued decreases in ocean pH mean that fluctuations in low pH will continue to reach levels below which sea urchin have experienced previously. For example, recent pH levels have dropped below 7.5, indicating that new extreme lows are beginning to occur (Chan et al. 2017). Therefore, we chose the extreme pH of 7.0 in order to understand how this species may respond to a novel, potentially impending, pH condition. Eggs for both the static and variable culturing treatments were fertilized in their respective static pH counterparts, with evenly pooled sperm from all males in 22-micron filtered, UV-sterilized seawater at 14°C with a salinity of 31ppt.

After verifying approximately >95% fertilization success (determined by the appearance of the fertilization envelope a few minutes after the addition of sperm), fertilized eggs were pooled across all females within each of the three static pH conditions (980,000 eggs per static pH condition). At 24 hours post fertilization ("day 1"), hatched blastula larvae were sampled from the three static pH pools ($N_{replicates} = 6$; 11,250 eggs per pH), and remaining blastulae were seeded into replicate culturing vessels $(N_{replicates} = 6; 11,250 \text{ eggs per } 3.7L \text{ vessel}; \text{ seeding density: } 3 \text{ larvae/mL})$ for each of the six different pH regimes: one control and two treatments that remained at a static pH (pH 8.1, pH 7.5, and pH 7.0) and 3 variable treatments that varied by 0.6 pH units over the course of 24 hours (pH 8.1 to 7.5, pH 7.8 to 7.2, and pH 7.3 to 6.7) (Fig. 1A), the daily change in pH that occurs during upwelling season in the CCME (Evans et al. 2013; Chan et al. 2017). It is important to note that the variable pH 8.1 treatment is meant to mimic the pH range commonly experienced in the CCME during upwelling (8.1 to 7.5) and has a mean pH of 7.8, whereas the edge and extreme varying pH treatments vary by the same amount but with the means matching their static edge and extreme pH treatment counterparts (pH 7.5 and pH 7.0).

Experimental system & water chemistry

Larvae were reared in a custom-designed, recirculating larval culturing system that allowed for continuous water movement set to a flow rate of 0.5 mL/sec from the top of each replicate culturing vessel exiting through a mesh covered cylinder in the bottom third of the vessel. The vessel design and flow rate were experimentally determined to minimize congregation of larvae at the outflow. For optimal water chemistry, 22-micron

filtered natural seawater was brought to UVM from the University of New Hampshire Coastal Marine Laboratory and UV-sterilized upon arrival. Seawater was maintained at a temperature of 14°C with a salinity of 31ppt. pH and temperature were measured by computer-monitored Hamilton Polilyte pH probes that were calibrated with three Thermo Scientific Orion pH buffers (4.01, 7.00, and 10.01). pH levels were controlled through communication between the pH probe, a computer system (RCK systems, San Diego, CA) and a solenoid valve that would release pure CO_2 gas through airstone bubblers measured and dosed every 10 seconds to maintain static or follow programmed variable pH conditions. Temperature was maintained by heat exchangers attached to the header tanks. CO₂ scrubbers were attached to the protein skimmers in the sump tanks for the pH 8.1 static (control) and variable treatment in order to help maintain control pH conditions for the former and to help pH increase when scheduled for the latter. Additional water chemistry measurements were taken on days 1, 4, and 7 post-fertilization (Table S1), with temperature, salinity, and pH measured around the same time of day and water collected and sealed for follow-up titration to determine total alkalinity (TA). TA was measured with a Mettler Toledo G10S Titrator, standardized to Andrew Dicksons' seawater standards (Dickson 2010). All of these water chemistry measurements were entered into the CO2Sys v.2.1 program to calculate pCO_2 (Pierrot et al. 2006). On days 3 and 5 postfertilization, recirculation was paused, and larvae were fed 1000 cells/mL each of Dunaliella spp. and Rhodomonas spp. (Brennan et al., 2019; Pespeni et al., 2013a) and allowed to feed for one hour before recirculation resumed. Larvae were reared in the culture vessels until 7 days old, the pluteus larval stage, at which point they were sampled for morphometric, survival, and genomic analyses (below).



Fig. 1: (A) pH measurements from the larval culturing system day 1 post-fertilization to end of the experiment (day 7 post-fertilization), recorded every 15 minutes from each of the 6 header tanks. The darker colors show the three static treatments (pH 8.1, 7.5, and 7.0) and the lighter colors show their varying counterparts. See legend in B. (B) Tukey boxplot of survival (percentage) after seven days of development in each treatment (n=12-18 survival estimates per pH condition). Letters above each boxplot indicate results from post hoc tests where different letters indicate significantly different groups. (C) Total body length (micrometers) of sampled surviving larvae (n=48-55 measurements per pH condition) after seven days of development.

Morphometrics & survival estimates

Seven-day old plutei were preserved in calcium carbonate-buffered formalin in seawater, and then photographed for morphometric analysis as in <u>Brennan et al. 2019 (n=48-55 per pH condition</u>). Total body length, represented as the midline body length extended to the top of the arms, was measured using *ImageJ* software (<u>Schneider et al., 2012</u>), calibrated with a stage micrometer. Data analysis was conducted in R (<u>R Core Team, 2015</u>) via a generalized linear model, with the family parameter set to 'gaussian' and pH as the main effect.

To estimate survival to day 7, culturing vessels were gently stirred, then three samples of 33mL were collected from each vessel and preserved in calcium carbonatebuffered formalin in seawater. Based on the starting density of 3 larvae/mL, 100% survival would yield ~100 larvae in 33 mL. Estimated percent survival was thus calculated based on the number of larvae counted over the expected 100 larvae for each of the three replicate samples per vessel. Larvae were considered to be alive if they had the pluteus larval developmental form, regardless of the number of arms, as low pH has been shown to cause developmental delay in urchins (Kurihara and Shirayama 2004). Abnormal larvae were minimal and not included in the alive count. Data analysis was conducted in R via a logistic regression using the *glm* function, with the response variable (survival) represented as two counts: 'alive' or 'dead' and pH treatment as the main effect. Post hoc comparisons were conducted using Tukey's Multiple Comparison of Means in the *multcomp* package (Hothorn et al., 2008).

DNA sequencing, processing, mapping, & SNP-calling

Four replicate pools of starting larval populations (day 1, n=11,250 larvae) per initial static pH condition (pH 8.1, pH 7.5, and pH 7.0) as well as the pools of surviving larvae (day 7, n= approximately 1000 to 3000 larvae) from each of the 6 culture vessels per pH condition, were collected and spun down to remove excess seawater, flash frozen in liquid nitrogen, and stored at -80 °C until extractions for genomic analyses. DNA was extracted with a Zymo ZR-Duet DNA/RNA MiniPrep Plus Kit (Zymo, Irvine, CA, USA). DNA was shipped over dry ice to Rapid Genomics (Gainesville, FL, USA) for library preparation and capture-sequencing with 46,316 custom 120bp probes, with two probes designed per gene based on the *S. purpuratus* genome v.3.1: one in an exonic region and the other in a putative regulatory region falling with 1000bp upstream of the transcription start site. Samples were barcoded, pooled, and then sequenced as 150bp paired-end reads on a single Illumina HiSeqX lane.

Raw DNA reads had Illumina adapters removed and were quality filtered with *trimmomatic-0.36* (Bolger et al. 2014) accepting final reads \geq 35bp in length. Filtered reads were mapped to the *S. purpuratus* genome v. 3.1 (build 7, echinobase.org) with BWA-MEM (Li 2013). SNPs were called using *Varscan* (Koboldt et al., 2012), resulting in 19,529,443 SNPs that were then further filtered in *R* (R Core Team, 2015) by depth (\geq 30, Ferretti et al. 2013), minor allele frequency (0.025), and a high coverage filter of 3 times the median coverage. This resulted in 54,427 high-quality SNPs for downstream analyses.

Principal Components Analysis was used to visualize variation across all identified SNPs among treatments and days. Significance among groups was tested using PERMANOVA implemented using the adonis function in the vegan package (v2.4-2). Cochran-Mantel-Haenszel (CMH) tests were used to identify specific loci with consistent changes in allele frequencies across replicates between pairs of groups with replicates randomly downsampled to four to have matched numbers of replicates for all contrasts. The CMH statistical test is an accurate method for identification of loci with changes in allele frequency in response to experimental selection (Vlachos et al. 2019). The test relies on sufficient replication; simulation studies show that the four replicates per condition should result in sufficient power to identify selected loci (Kofler and Schlötterer 2014). We tested for differences in allele frequency relative to both the day 1, pH 8.1 samples and the day 7, pH 8.1 static samples. Results from all contrasts are presented in Table S2. We focus downstream analyses using the contrasts to day 7, pH 8.1 static samples to control for potential changes in allele frequency due to lab adaptation. Linkage disequilibrium decays quickly in S. purpuratus, over a few hundred base pairs (Brennan et al. 2019). As such, we considered loci associated with each gene region (combining coding and regulatory regions for a given gene) to be independent, resulting in 4,548 independent regions. Given the low linkage disequilibrium, the number of independent regions represents the number of independent tests. We thus used a stringent Bonferroni correction to adjust for multiple testing, where 0.05/4548 resulted in a *p*-value threshold of 1.1×10^{-5} . To test for correlations of changes in allele frequency within and between the low and extreme static and variable pH conditions, we used Pearson's Correlation. Finally, loci were categorized as exonic (synonymous, nonsynonymous), intronic, promoter, or intergenic and linked to genes using SnpEff (Cingolani et al. 2012). χ^2 tests were used to test if any group contained significantly more loci responding to selection than expected by chance. Loci significantly changing in allele frequency for each treatment were tested for functional enrichment of gene ontology (GO) categories. GO terms for annotated genes were downloaded from EchinoBase (www.echinobase.org). TOPGO v. 2.36.0 was used to test for significant enrichment of function categories using the weight algorithm and limited to terms that had at least 5 annotated genes (Alexa et al. 2006). Similarity among significant GO terms (P < 0.05) was calculated with GOSemSim in R using Wang's method, which calculates similarity based on the topology of the GO graph structure (Yu et al. 2010). This similarity measure was converted to a dissimilarity matrix, hierarchically clustered, and

plotted using ggdendro in R (de Vries and Ripley 2016).

Code and details to reproduce all analyses are available at: https://github.com/PespeniLab/spoa static vs variable.

Results

Survival and morphometrics

pH condition had a significant effect on both larval survival and body size where lower pH, generally, resulted in higher mortality and smaller body size. Larval survival (n=12-18 samples per pH treatment) decreased by 20% after 7 days in the static pH 7.5 (edge) and pH 7.0 (extreme) conditions, compared to the static control, pH 8.1 (Fig. 1B, p < 0.001). No difference was found in survival between the edge and extreme static conditions (P = 0.44). Under the pH 7.5 selection regime, variability improved survival relative to static conditions such that survival was comparable to control static conditions (pH 8.1). A similar mitigation was observed for pH 7.0 under variable conditions, which was increased compared to its static counterpart (Fig. 1B, P < 0.001), though survival did not improve to match control conditions. In agreement with survival results, total body length of 7-day plutei showed a similar reduction as pH decreased, from a mean size of 319 micrometers to 230 and 175 micrometers for the static pH 7.5 and 7.0 conditions, respectively (Fig. 1C, P < 0.001). However, variable conditions did not result in the same pattern of recovery. In contrast, variable conditions had either similar sized (pH 7.5, mean = 225 micrometers) or smaller individuals (pH 8.1 and 7.0, mean = 268 and 151 micrometers, respectively) than their static counterparts (Fig. 1C, P < 0.001).

Genomic Analyses

Genome-wide allele frequency estimates across all 54,427 high quality SNPs identified from pooled, capture-sequenced genomic DNA showed low variation among replicates within each treatment group using Principal Components Analysis (PCA, Fig. 2A). Samples clustered separately according to mean pH and day of sampling (PERMANOVA, P < 0.05), where samples moved sequentially further in PC space from the day 1, pH 8.1 control as the low pH condition intensified. Accordingly, the largest differences observed were between day 1 control static condition and day 7, pH 7.0 (Fig. 2A). Static and variable treatments within a pH overlapped in their distributions except for pH 7.0 (PERMANOVA, P = 0.04), suggesting stronger selection in response to mean pH rather than variability. Despite broad differences in allele frequencies and high levels of mortality, no differences in estimates of nucleotide diversity were observed among day or treatment groups (Table S3).



Fig. 2: (A) Principal component analysis of the 54,427 high quality SNPs for the six pH conditions and two sampled time periods. Color indicates pH condition in the bottom portion of the legend key, where dark colors are static and light are variable. Shapes

(circle or triangle) denote which day post-fertilization; circles are samples from day one of development and triangles from samples after seven days of development. (B) *Venn diagram* of loci with significant changes in allele frequency in response to selection at pH 7.5 and 7.0 for static and variable conditions (compared to day 7, static pH 8.1).



Fig. 3: Number of loci significantly changing in frequency in different regions of the genome. P-values in the top right corners indicate corrected significance from χ^2 tests where values are FDR corrected to account for multiple testing. Within each panel, expected values are from χ^2 tests and observed values are the actual counts of significant loci. Error bars for observed values represent the 95% confidence interval calculated with the Clopper-Pearson method (Clopper and Pearson 1934). Asterisks indicate groups with a significantly greater number of loci than expected by chance (corresponding grey bar), as determined by binomial post-hoc tests. Note that we have chosen to collapse synonymous and non-synonymous loci in exonic regions as linkage disequilibrium extends at least 100 bp (Brennan et al. 2019) inhibiting identification of specific loci

targeted by selection. See Fig. S1 for these loci separated.

CMH tests identified loci consistently responding to pH selection after 7 days of development compared to the day 7, static pH 8.1. Across the treatments, decreasing pH resulted in a greater number of loci with significant changes in allele frequencies relative to the static pH 8.1 control, a pattern that agrees with the increasing distance in principle component space as pH decreases. We observed limited changes in allele frequency between the static pH 8.1 and variable (mean pH 7.8) conditions, with only 6 divergent loci ($P < 1.1 \times 10^{-5}$). More divergence was observed in pH 7.5 with 24 and 34 pHselected loci for static and variable treatments, respectively (Fig. 2B). However, of these loci, only 4 were overlapping (7.4% of pH 7.5 selected loci). We observed the largest degree of allelic divergence from the control at pH 7.0 with 602 and 253 significant loci in the respective static and variable treatments and 106 overlapping variants between them (14.2% of the pH 7.0 selected loci; Fig. 2B). Between the pH treatments, we observed little overlap in responsive loci (1% of static selected loci; 0.3% of variable selected loci; Fig. 2B), indicating the genetic responses to different degrees of pH stress were largely unique. Finally, we found that loci in exonic regions were more likely to be targets of selection than loci in promoter, intronic, or intergenic regions ($\chi^{2df=3} = 100$, P < 0.001; Fig. 3). This pattern was driven by pH 7.0 variable and static (P < 0.001) and pH 7.5 variable (P = 0.055), though both pH 7.5 static and pH 8.1 follow a similar trend (Fig. 3).



Fig. 4: Correlation between loci showing significant changes in allele frequency ($P < 1.1 \times 10-5$) in response to selection in different treatments. Change in allele frequency for all plots are relative to frequencies of day 7, static pH 8.1 static at day 7. Color and shape indicate each treatment while loci significant for both treatments ('overlapping') in each panel are represented by yellow diamonds. (A) within pH 7.0, static versus variable (r = 0.54). (B) within pH 7.5, static versus variable (r = 0.21). (C) within static, pH 7.0 versus pH 7.5 (r = 0.47). (D) within variable, pH 7.0 versus pH 7.5 (r = 0.25).

We observed correlated changes in allele frequencies within and between pH conditions (Fig. 4). Within pH 7.0, changes in allele frequency in response to static and variable conditions were strongly correlated (Pearson's r = 0.54; Fig. 4A). Conversely, static and variable responses within pH 7.5 showed a weaker correlation (r = 0.21; Fig. 4B). Comparing static and variable responses between pH 7.0 and 7.5 revealed different patterns. Changes in frequency in response to static pH were strongly correlated between pH 7.5 and 7.0 (r = 0.47; Fig. 4C) but the correlation between the variable treatments was much weaker (r = 0.25; Fig 4D).



Fig. 5: Gene ontology enrichment results for biological processes. Checks indicate enrichment ($P_{adj} < 0.05$) for significant changes in allele frequency for loci in the category (row) in response to the treatment indicated (column). GO terms are clustered by similarity according to their topology in the GO graph structure. See methods for details.

Functional enrichment analyses revealed unique and shared biological processes subject to selection among the experimental groups (Fig. 5). See Figure S1 and tables S4-S8 for enrichment results for cellular components, molecular function, and for day 1 (D1) vs. day 7 (D7) comparisons. The variable pH 8.1 treatment (mean pH 7.8) was enriched for one category related to hormone-mediated signaling. In response to pH 7.5 conditions, the majority of categories enriched for changes in allele frequency were involved in the biosynthesis, metabolism, transportation, or modification of lipids. Three categories were shared between the pH 7.5 static and variable conditions, all related to phosphatidylinositol metabolism, signaling, and phosphorylation. In response to extreme pH 7.0 conditions, allele frequency changes were also enriched for lipid and phospholipid processes. However, there were several processes unique to pH 7.0, including transmembrane and mitochondrial processes, and RNA processing. Though the pH 7.0 static and variable treatments had overlapping and correlated genetic responses (Figs. 2B and 4A), only one GO term overlapped between the two treatments: cellular protein complex disassembly. In static and variable pH 7.0 conditions, selection disproportionately targeted exonic regions. Consistent with overall enrichment in these conditions, selected exonic loci in static pH 7.0 included immune responses (Table S9) while exonic loci in variable pH 7.0 were enriched for vesicle transport and metabolic processes (Table S10). Static conditions at different pH shared enrichment for immune responses, while the variable conditions shared no common functional categories. Interestingly, there were similar numbers of enriched categories for surviving pH 7.0 and pH 7.5, despite the 10-fold greater number of loci changing in allele frequency in response to the more extreme pH environment.

Discussion

We find that as seawater acidification increases in intensity, both the physiological (Fig. 1) and genomic (Figs. 2-4) consequences become more extreme. In particular, when pH conditions extend beyond what is currently experienced in nature, the adverse effects on the survival and growth of developing sea urchin larvae are most pronounced. We demonstrate that while fluctuating pH conditions can reduce mortality relative to static conditions, the same effect is not seen for body size. Yet, we simultaneously show that adaptive genetic variation for extreme and fluctuating pH is present in wild populations. Loci targeted by selection were disproportionately found in genes related to lipid metabolism (pH 7.5) and membrane function (pH 7.0) and in exonic regions, indicating that changes to protein function enabled survival particularly in extreme pH 7.0 conditions. Importantly, the genetic responses to pH 7.5 (edge) and pH 7.0 (extreme) conditions were unique, indicating that the physiological processes enabling persistence are similarly unique. This suggests that changes along the logarithmic pH scale do not elicit a linear physiological response. Further, the distinct responses suggest that adaptation to gradual decreases in average pH may not facilitate persistence to extreme pH conditions, a hypothesis that warrants further investigation.

Tradeoff between survival and growth

S. purpuratus relies on amorphous calcium carbonate to lay down its skeleton during development (<u>Addadi et al. 2003</u>; Politi et al. 2008; <u>Vidavsky et al. 2014</u>) and, as pH decreases, the increase in energetic demands to develop skeletal structure leads to developmental abnormalities and higher mortality rates (<u>Stumpp et al. 2012</u>; <u>Pan et al.</u>

2015). Accordingly, the increased mortality and decreased size with increasing pH stress observed here (Fig. 1A) match expectations from previous work that has also shown reductions in size and survival under decreased pH conditions (Kurihara 2008; Ries et al. 2009; Stumpp et al. 2012). However, we find that fluctuating pH drives higher survival, but not body size (Fig. 1B,C), partially matching our predictions. Previous work has identified both positive (Dufault et al. 2012) and negative (Mangan et al. 2017; Onitsuka et al. 2018; Chan and Tong 2020) impacts of pH fluctuations on survival and growth. For purple sea urchin, fluctuations in pH may serve as a buffer, where periodic exposure to less stressful pH conditions enable more typical development that increases survival. Alternatively, a fluctuating environment is regularly experienced during the spawning season (Miller and Emlet 1997) and may represent the conditions to which individuals are adapted and thus best able to survive.

Our results suggest that, for early-in-development larvae, smaller body size may enable increased survival under low pH conditions; fluctuating conditions increase survival but decrease body size relative to static treatments (Fig. 1B,C). Decreasing pH consistently results in smaller larvae through development for S. purpuratus, among other sea urchin species as well (Yu et al. 2011; Suwa et al. 2013; Pespeni et al. 2013a; Brennan et al. 2019), but the adaptive significance of this change has been unclear. Larvae under low pH dedicate much of their energy to acid-base balance at the cost of growth (Stumpp et al. 2012; Pan et al. 2015). We hypothesize that smaller size and lower growth under low pH conditions decreases the total energy budget and allows for increased survival. Further, the success of metamorphosis is dependent on lipid energy reserves, which may be similarly increased by reducing size during development (Sewell 2005; Byrne et al. 2008). This is further corroborated by our GO enrichment results, which show increasing enrichment for genes related to lipid metabolism as pH decreases (Fig. 5). While small size during development may increase early survival or success of metamorphosis, it could also reduce overall population fitness. For example small larvae experience higher predation rates (Allen 2008), and may develop into smaller, less fecund adults (Dupont et al. 2013). Future work should address the consequences of reduced body size on energetic demands, survival, metamorphosis, and adult fitness in *S. purpuratus*.

Surviving in pH 7.5 (edge) versus pH 7.0 (extreme) conditions

Previous work has shown that populations of *S. purpuratus* harbor sufficient standing genetic variation to respond to static pH levels within (pH 7.8) and at the edge (pH 7.5) of their current range (Pespeni et al. 2013a; Brennan et al. 2019). Here, we demonstrate how adaptation may proceed as pH begins to extend beyond what is currently experienced in the wild. We find that responses to pH conditions beyond the natural range (pH 7.0) are not merely more extreme changes in allele frequency at the loci underlying adaptation to pH conditions within the natural range of pH variability (pH 7.5). Rather, selection targets a unique set of loci (Fig. 4B). Under extreme static pH conditions, we find selected loci in gene functions related to immune response, cell adhesion, mitochondrial gene expression, and lipid metabolism (Fig. 5). Immune response under extremely stressful conditions is not unexpected (Bibby et al. 2008; Brothers et al. 2016), especially in such an energetically-costly environment as low pH for calcifying larvae that need to maintain acid-base physiology (Stumpp et al. 2012; Pan

<u>et al. 2015</u>). Indeed, perhaps the enrichment for mitochondrial gene expression is being used by larvae under extreme conditions for energy allocation in order to pay the cost of surviving and maintaining cellular processes in novel pH conditions.

Loci under selection at the edge of the natural pH range (pH 7.5) are unique from those responding to pH 7.0 but overlap in a number of gene functions including immune response and lipid metabolism. Lipids are important for cellular membrane structure and serve as critical energy stores for calcifying marine invertebrates (Sewell 2005; Schoepf et al. 2013). Indeed, gene functions related to lipid metabolic processes were the only ones enriched to varying degrees across all of the edge and extreme pH treatments (Fig. 5). Lipid energy allocation in sea urchins is essential for successful metamorphosis, as metamorphosing larvae spend weeks nutritionally reliant on the lipid stores from the pluteus larval stage (Sewell 2005). Thus, high enrichment for lipid metabolism may confer a selective advantage for larvae subjected to low pH levels at the edge of their range. While we see no difference in survival between pH 7.5 and pH 7.0 static conditions (Fig 1B), the noticeably lower enrichment of lipid metabolism in static pH 7.0 conditions compared to static pH 7.5 suggests that genetic variation in lipid metabolism genes was critical for survival in pH 7.5. Without selection for these alleles, larvae in pH 7.0 may accrue a greater metabolic cost of pH stress resulting in negative carryover effects to later life history stages, such as metamorphosis and juvenile survival. Finally, for static pH 7.5, we observe unique enrichment for gene functions related to chemotaxis. Chemotaxis is especially important for phagocytes in sea urchin, which make up the majority of the coelomocytes, the first line of cellular defense in the sea urchin innate immune system (Smith et al. 2006), and have been shown to be negatively impacted by

acidification (Brothers et al. 2016).

Surviving static versus fluctuating pH conditions

Our findings suggest that genetic responses to variability are similar to static conditions (Fig 4); response to selection is correlated between the regimes, and related genetic mechanisms are used to respond to static and variable conditions. Considering the high amount of environmental variation in the CCME, along with the high levels of standing genetic variation found in this species (Sodergren et al. 2006; Pespeni and Palumbi 2013c), one explanation is that previous selection in fluctuating pH has led to genetic variation that is adaptive in pH-variable environments and similar mechanisms are used to survive static conditions with the same mean pH. We find enrichment in pH 7.0 (extreme) variable conditions for loci in genes related to lipid metabolism, ribosomal and RNA structure, and mitochondria structure and transport (Fig. 5). Previous OA research in copepods has shown multigenerational selection on ribosomal structure and oxidative phosphorylation, a key metabolic process that produces ATP and occurs within the mitochondria (De Wit et al. 2016).

The extreme variable pH 7.0 treatment exhibited a trade-off between survival and growth, with high survival in the pH 7.0 variable conditions compared to the pH 7.0 static, but at the cost of reduced size in surviving larvae. Integrating organismal and gene function results, purple urchin larvae may focus on energy production and allocation in order to survive extreme fluctuating pH conditions, putting more energy into survival over development. Differential energy allocation has been found previously in *S. purpuratus* (Pan et al. 2015) and multiple calcifying marine gastropod species, which use

'dwarfing' as an adaptive strategy to tolerate low pH (Garilli et al. 2015). Metabolomic work in another marine calcifier, *Pocillopora damicornis* coral, conveys the importance of cellular structure and maintenance for survival under pH stress, and suggests the energetic expense of this could come at a cost on growth (Sogin et al. 2016). Overall, while there is limited overlap in the specific functional categories enriched for static and variable pH conditions, enriched functions serve a shared purpose: maintenance of structural integrity and regulation of metabolism to simultaneously manage development, growth, and survival as pH decreases and/or fluctuates.

Conclusion

Understanding the genetic and phenotypic responses of organisms to their environment is pertinent, especially in a rapidly changing climate characterized by increases in extreme and variable conditions. Here, we show that purple sea urchins have greater adaptive potential than previously thought, with genetic variation available to respond to extreme static and variable pH conditions. pH variability, while leading to survivors with smaller bodies, did increase larval survival. High levels of standing genetic variation, coupled with natural variation in pH conditions across space and time, appear to promote the maintenance of adaptive potential for purple sea urchin populations. Future studies, however, should explore carryover effects to later life history stages and across generations and test the hypothesis that these factors promote adaptive potential to global change conditions in a broader phylogenetic framework. Ultimately, environmental change may be buffered against for species that have sufficient genetic variation for responding to such change.

Funding

This work was supported by the National Science Foundation (NSF) Graduate Research Fellowship Program DGE-1451866 (to A.D.G.) and NSF grants OCE-1559075 and OIA-1736253 (to M.H.P.).

Acknowledgements

We thank Pete Halmay and Patrick Leahy for urchin collections, UNH Coastal Marine Lab for providing filtered seawater, Michael Paquette and Aqua Logic for the experimental system design and related troubleshooting, Dr. Justin McAlister and Lauren Ashlock for providing the algal species used to feed larvae, and the Pespeni Lab undergraduate cohort that greatly assisted with preparing and running the experiment (Emily Shore, Chelsea Darwin, Malcolm Hughes, Erika Petterssen, Seth Lowell, Katherine Helmer, and Grace King). We also thank Leigh Sweet for helping with the seawater collection and helpful troubleshooting discussions, several UVM faculty for helpful discussions about experimental design and analysis (Dr. Brent Lockwood, Dr. Steve Keller, and Dr. Alicia Ebert), and Mike Austin and UVM's ETS for server space and maintenance. **Supplemental Figures and Tables**



Fig. S1: Number of loci significantly changing in frequency in different regions of the genome, where exonic regions are split into synonymous and non-synonymous sites. P-values in the top right corners indicate corrected significance from χ^2 tests where values are FDR corrected to account for multiple testing. Within each panel, expected values are from χ^2 tests and observed values are the actual counts of significant loci. Error bars for observed values represent the 95% confidence interval calculated with the Clopper-Pearson method. Asterisks indicate groups with a significantly greater number of loci than expected by chance (corresponding grey bar), as determined by binomial post-hoc tests.

Table S1: Water chemistry data recorded on days 1, 4, and 7 post-fertilization. A water sample was taken from each of the six pH condition header tanks, with temperature, pH, and salinity recorded immediately while total alkalinity (TA) was measured via titration and then pCO2 calculated from CO2Sys.

Day	Treatment	рН	Temp (℃)	Salinity (ppt)	TA (umol/kg SW)	pCO2 (uatm)
1	8.1S	8.10	14.1	30.5	2085.372	317.3
	8.1V	8.09	14.5	30.5	2019.325	315.3
	7.5S	7.50	14.2	30.5	2078.033	1444.0
	7.5V	7.51	14.0	30.5	2010.763	1362.0
	7.0S	7.00	14.2	30.5	2093.114	4760.7
	7.0V	7.01	14.1	30.5	2073.548	4604.1
4	8.1S	8.10	13.9	30.5	1976.260	300.1
	8.1V	7.70	13.8	30.5	2025.208	861.2
	7.5S	7.50	14.2	30.5	2025.269	1406.9
	7.5V	7.23	14.1	30.5	1998.532	2644.4
	7.0S	7.00	13.8	30.0	2016.642	4588.0
	7.0V	6.77	13.9	30.5	2063.356	8007.5
7	8.1S	8.10	13.9	31.5	2066.227	311.0
	8.1V	7.62	13.9	30.5	2056.235	1065.5
	7.5S	7.50	14.0	31.5	2055.902	1417.8
	7.5V	7.30	13.9	31.5	2093.933	2331.1
	7.0S	6.95	14.0	30.5	2038.123	5203.4
	7.0V	6.82	14.2	32.5	2083.740	7140.6

Table S2: Number of significant loci from the CMH tests. Comparisons were done between all of the different days and pH conditions, comparing to both the day 1, 8.1S (top row) and day 7, 8.1S (bottom row).

	D7 8.1 S	D7 8.1 V	D7 7.5 S	D7 7.5 V	D7 7.0 S	D7 7.0 V	D1 7.5 S	D1 7.0 S
D1 8.1 S	139	152	73	169	1057	352	70	772
D7 8.1 S		6	24	34	602	253		

Table S3: Nucleotide diversity results for all of the different days and pH conditions.

Treatment	D1 8.1S	D7 8.1S	D7 7.5S	D7 7.5V	D7 7.0S	D7 7.0V	D7 8.1V
Nucleotide Diversity	0.0138	0.0143	0.0136	0.0140	0.0139	0.0134	0.0143

Table S4: Gene ontology enrichment results for (A) molecular functions and (B)cellular components. Checks indicate enrichment ($P_{adj} < 0.05$) for significant changes in

allele frequency for loci in the category (row) in response to the treatment indicated (column). GO terms are clustered by similarity according to their topology in the GO graph structure.

(A)



*Note: Supplemental Tables S5-S10 are text files that can be found online with the corresponding publication of this thesis (citation found on page ii of this thesis).

BIBLIOGRAPHY

Addadi L, Raz S, Weiner S. 2003. Taking Advantage of Disorder: Amorphous Calcium Carbonate and Its Roles in Biomineralization. Adv Mater 15:959–70.

Alexa A, Rahnenfuhrer J. 2006. TopGO: enrichment analysis for gene ontology. R package version.

Allen JD. 2008. Size-specific predation on marine invertebrate larvae. Biol Bull 214:42–49.

Barrett RDH, Laurent S, Mallarino R, Pfeifer SP, Xu CCY, Foll M, Wakamatsu K, Duke-Cohan JS, Jensen JD, Hoekstra HE. 2019. Linking a mutation to survival in wild mice. Science 363:499–504.

Beardmore JA, Levine L. 1963. Fitness and Environmental Variation. 1. A Study of Some Polymorphic Populations of *Drosophila pseudoobscura*. Evolution 17:121–29.

Bibby R, Widdicombe S, Parry H, Spicer J, Pipe R. 2008. Effects of ocean acidification on the immune response of the blue mussel *Mytilus edulis*. Aquat Biol 2:67–74.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–20.

Bozinovic F, Bastías DA, Boher F, Clavijo-Baquet S, Estay SA, Angilletta MJ Jr. 2011. The mean and variance of environmental temperature interact to determine physiological tolerance and fitness. Physiol Biochem Zool 84:543–52.

Brennan RS, Garrett AD, Huber KE, Hargarten H, Pespeni MH. 2019. Rare genetic variation and balanced polymorphisms are important for survival in global change conditions. Proc Biol Sci 286:20190943.

Brothers CJ, Harianto J, McClintock JB, Byrne M. 2016. Sea urchins in a high-CO2 world: the influence of acclimation on the immune response to ocean warming and acidification. Proc Biol Sci 283.

Byrne M, Prowse TAA, Sewell MA, Dworjanyn S, Williamson JE, Vaïtilingon D. 2008. Maternal provisioning for larvae and larval provisioning for juveniles in the toxopneustid sea urchin *Tripneustes gratilla*. Mar Biol 155:473–82.

Byrne M, Lamare M, Winter D, Dworjanyn SA, Uthicke S. 2013. The stunting effect of a high CO2 ocean on calcification and development in sea urchin larvae, a synthesis from the tropics to the poles. Philos Trans R Soc Lond B Biol Sci 368:20120439.

Calosi P, De Wit P, Thor P, Dupont S. 2016. Will life find a way? Evolution of marine species under global change. Evol Appl 9:1035–42.

Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards SV. 2017. Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole lizard. Science 357:495–98.

Carlson SM, Cunningham CJ, Westley PAH. 2014. Evolutionary rescue in a changing world. Trends Ecol Evol 29:521–30.

Chan F, Barth JA, Blanchette CA, Byrne RH, Chavez F, Cheriton O, Feely RA, Friederich G, Gaylord B, Gouhier T, Hacker S, Hill T, Hofmann G, McManus MA, Menge BA, Nielsen KJ, Russell A, Sanford E, Sevadjian J, Washburn L. 2017. Persistent spatial structuring of coastal ocean acidification in the California Current System. Sci Rep 7:2526.

Chan KYK, Tong CSD. 2020. Temporal variability modulates pH impact on larval sea urchin development: Themed Issue Article: Biomechanics and Climate Change. Conserv Physiol 8:coaa008.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly 6:80–92.

de Vries A, Ripley BD. 2016. ggdendro: Create Dendrograms and Tree Diagrams Using 'ggplot2'. R package version 0 1-20.

De Wit P, Dupont S, Thor P. 2016. Selection on oxidative phosphorylation and ribosomal structure as a multigenerational response to ocean acidification in the common copepod *Pseudocalanus acuspes*. Evol Appl 9:1112–23.

Dickson AG. 2010. The carbon dioxide system in seawater: equilibrium chemistry and measurements. Guide to best practices for ocean acidification research and data reporting 1:17–40.

Doney SC, Fabry VJ, Feely RA, Kleypas JA. 2009. Ocean acidification: the other CO2 problem. Ann Rev Mar Sci 1:169–92.

Dufault AM, Cumbo VR, Fan T-Y, Edmunds PJ. 2012. Effects of diurnally oscillating pCO₂ on the calcification and survival of coral recruits. Proc Biol Sci 279:2951–58.

Dupont S, Dorey N, Stumpp M, Melzner F, Thorndyke M. 2013. Long-term and translife-cycle effects of exposure to ocean acidification in the green sea urchin *Strongylocentrotus droebachiensis*. Mar Biol 160:1835–43. Easterling DR, Meehl GA, Parmesan C, Changnon SA, Karl TR, Mearns LO. 2000. Climate extremes: observations, modeling, and impacts. Science 289:2068–74.

Esbaugh AJ, Heuer R, Grosell M. 2012. Impacts of ocean acidification on respiratory gas exchange and acid–base balance in a marine teleost, *Opsanus beta*. J Comp Physiol B 182:921–34.

Estay SA, Clavijo-Baquet S, Lima M, Bozinovic F. 2011. Beyond average: an experimental test of temperature variability on the population dynamics of *Tribolium confusum*. Population Ecology.

Evans TG, Chan F, Menge BA, Hofmann GE. 2013. Transcriptomic responses to ocean acidification in larval sea urchins from a naturally variable pH environment. Mol Ecol 22:1609–25.

Fabry VJ, Seibel BA, Feely RA, Orr JC. 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. ICES J Mar Sci 65:414–32.

Ferretti L, Ramos-Onsins SE, Pérez-Enciso M. 2013. Population genomics from pool sequencing. Mol Ecol 22:5561–76.

Flanagan SP, Forester BR, Latch EK, Aitken SN, Hoban S. 2018. Guidelines for planning genomic assessment and monitoring of locally adaptive variation to inform species conservation. Evol Appl 11:1035–52.

Folguera G, Bastías DA, Caers J, Rojas JM, Piulachs M-D, Bellés X, Bozinovic F. 2011. An experimental test of the role of environmental temperature variability on ectotherm molecular, physiological and life-history traits: Implications for global warming. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology.

Frieder CA, Gonzalez JP, Bockmon EE, Navarro MO, Levin LA. 2014. Can variable pH and low oxygen moderate ocean acidification outcomes for mussel larvae? Glob Chang Biol 20:754–64.

Garilli V, Rodolfo-Metalpa R, Scuderi D, Brusca L, Parrinello D, Rastrick SPS, Foggo A, Twitchett RJ, Hall-Spencer JM, Milazzo M. 2015. Physiological advantages of dwarfing in surviving extinctions in high-CO2 oceans. Nat Clim Chang 5:678–82.

Gompert Z, Comeault AA, Farkas TE, Feder JL, Parchman TL, Buerkle CA, Nosil P. 2014. Experimental evidence for ecological selection on genome variation in the wild. Ecol Lett 17:369–79.

Hoffmann AA, Sgrò CM. 2011. Climate change and evolutionary adaptation. Nature 470:479–85.

Hofmann GE, Smith JE, Johnson KS, Send U, Levin LA, Micheli F, Paytan A, Price NN,

Peterson B, Takeshita Y, Matson PG, Crook ED, Kroeker KJ, Gambi MC, Rivest EB, Frieder CA, Yu PC, Martz TR. 2011. High-frequency dynamics of ocean pH: a multiecosystem comparison. PLoS One 6:e28983.

Hothorn T, Bretz F, Westfall P. 2008. Simultaneous inference in general parametric models. Biom J 50:346–63.

IPCC. 2019. Technical Summary [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, E. Poloczanska, K. Mintenbeck, M. Tignor, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. In press.

Jarrold MD, Humphrey C, McCormick MI, Munday PL. 2017. Diel CO2 cycles reduce severity of behavioural abnormalities in coral reef fish under ocean acidification. Sci Rep 7:10153.

Katz RW, Brown BG. 1992. Extreme events in a changing climate: Variability is more important than averages. Clim Change 21:289–302.

Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L, Wilson RK. 2012. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. Genome Res 22:568–76.

Kroeker KJ, Kordas RL, Crim RN, Singh GG. 2010. Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. Ecol Lett 13:1419–34.

Kurihara H, Shirayama Y. 2004. Effects of increased atmospheric CO2 on sea urchin early development. Mar Ecol Prog Ser 274:161–69.

Kurihara H. 2008. Effects of CO2-driven ocean acidification on the early developmental stages of invertebrates. Mar Ecol Prog Ser 373:275–84.

Kwiatkowski L, Orr JC. 2018. Diverging seasonal extremes for ocean acidification during the twenty-first century. Nat Clim Chang 8:141–45.

Lande R, Shannon S. 1996. The role of genetic variation in adaptation and population persistence in a changing environment. Evolution 50:434–37.

Leahy PS. 1986. Laboratory culture of Strongylocentrotus purpuratus adults, embryos, and larvae. Methods Cell Biol 27:1–13.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv [q-bioGN].

Li F, Wu Y, Hutchins DA, Fu F, Gao K. 2016. Physiological responses of coastal and oceanic diatoms to diurnal fluctuations in seawater carbonate chemistry under two CO 2 concentrations. Biogeosciences 13.

Long T. 1970. Genetic effects of fluctuating temperature in populations of *Drosophila melanogaster*. Genetics 66:401–16.

Lynch M, Walsh B, Others. 1998. Genetics and analysis of quantitative traits Sinauer Sunderland, MA.

Mangan S, Urbina MA, Findlay HS, Wilson RW, Lewis C. 2017. Fluctuating seawater pH/pCO₂ regimes are more energetically expensive than static pH/ pCO₂ levels in the mussel *Mytilus edulis*. Proc Biol Sci 284.

Marshall DJ, Morgan SG. 2011. Ecological and evolutionary consequences of linked lifehistory stages in the sea. Curr Biol 21:R718–25.

Martins NE, Faria VG, Nolte V, Schlötterer C, Teixeira L, Sucena É, Magalhães S. 2014. Host adaptation to viruses relies on few genes with different cross-resistance properties. Proc Natl Acad Sci U S A 111:5938–43.

McNeil BI, Sasse TP. 2016. Future ocean hypercapnia driven by anthropogenic amplification of the natural pCO₂ cycle. Nature 529:383–86.

Miller BA, Emlet RB. 1997. Influence of nearshore hydrodynamics on larval abundance and settlement of sea urchins *Strongylocentrotus franciscanus* and *S. purpuratus* in the Oregon upwelling zone. Vol 148:83–94.

Nielsen R, Slatkin M. 2013. An introduction to population genetics. Sunderland, MA: Sinauer Associates [Google Scholar].

Onitsuka T, Takami H, Muraoka D, Matsumoto Y, Nakatsubo A, Kimura R, Ono T, Nojiri Y. 2018. Effects of ocean acidification with pCO₂ diurnal fluctuations on survival and larval shell formation of Ezo abalone, *Haliotis discus hannai*. Mar Environ Res 134:28–36.

Palumbi SR, Wilson AC. 1990. Mitochondrial DNA diversity in the sea urchins Strongylocentrotus purpuratus and S. droebachiensis. Evolution 44:403–15.

Pan T-CF, Applebaum SL, Manahan DT. 2015. Experimental ocean acidification alters the allocation of metabolic energy. Proc Natl Acad Sci U S A 112:4696–4701.

Pearse JS. 2006. Ecological role of purple sea urchins. Science 314:940–41.

Pespeni MH, Sanford E, Gaylord B, Hill TM, Hosfelt JD, Jaris HK, LaVigne M, Lenz EA, Russell AD, Young MK, Palumbi SR. 2013a. Evolutionary change during

experimental ocean acidification. Proc Natl Acad Sci U S A 110:6937-42.

Pespeni MH, Chan F, Menge BA, Palumbi SR. 2013b. Signs of adaptation to local pH conditions across an environmental mosaic in the California Current Ecosystem. Integr Comp Biol 53:857–70.

Pespeni MH, Palumbi SR. 2013c. Signals of selection in outlier loci in a widely dispersing species across an environmental mosaic. Mol Ecol 22:3580–97.

Pierrot D, Lewis E, Wallace DWR, Others. 2006. CO2SYS DOS Program developed for CO2 system calculations. ORNL/CDIAC-105 Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, TN.

Politi Y, Metzler RA, Abrecht M, Gilbert B, Wilt FH, Sagi I, Addadi L, Weiner S, Gilbert PUPA. 2008. Transformation mechanism of amorphous calcium carbonate into calcite in the sea urchin larval spicule. Proc Natl Acad Sci U S A 105:17362–66.

Price NN, Martz TR, Brainard RE, Smith JE. 2012. Diel variability in seawater pH relates to calcification and benthic community structure on coral reefs. PLoS One 7:e43843.

Rackley SA. 2017. Carbon Capture and Storage Butterworth-Heinemann.

R Core Team. 2015. R: A language and environment for statistical computing. *R Foundation for Statistical Computing*, Vienna, Austria. URL <u>http://www.R-project.org</u>.

Rahmstorf S, Coumou D. 2011. Increase of extreme events in a warming world. Proc Natl Acad Sci U S A 108:17905–9.

Ries JB, Cohen AL, McCorkle DC. 2009. Marine calcifiers exhibit mixed responses to CO2-induced ocean acidification. Geology 37:1131–34.

Rogers-Bennett L. 2007. Chapter 19 The ecology of Strongylocentrotus franciscanus and Strongylocentrotus purpuratus. Developments in Aquaculture and Fisheries Science.

Samuelsson T, Olsson M. 1990. Transfer RNA Pseudouridine Synthases in *Saccharomyces cerevisiae*. Plan Perspect 8782:8787.

Schlötterer C, Kofler R, Versace E, Tobler R, Franssen SU. 2016. Combining experimental evolution with next-generation sequencing: a powerful tool to study adaptation from standing genetic variation. Heredity 116:248.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. Nat Methods 9:671–75.

Schoepf V, Grottoli AG, Warner ME, Cai W-J, Melman TF, Hoadley KD, Pettay DT, Hu

X, Li Q, Xu H, Wang Y, Matsui Y, Baumann JH. 2013. Coral energy reserves and calcification in a high-CO2 world at two temperatures. PLoS One 8:e75049.

Sewell MA. 2005. Utilization of lipids during early development of the sea urchin *Evechinus chloroticus*. Mar Ecol Prog Ser 304:133–42.

Shama LNS. 2017. The mean and variance of climate change in the oceans: hidden evolutionary potential under stochastic environmental variability in marine sticklebacks. Sci Rep 7:8889.

Smith LC, Rast JP, Brockton V, Terwilliger DP, Nair SV, Buckley KM, Majeske AJ. 2006. The sea urchin immune system. Invertebrate Surviv J 3:25–39.

Sodergren E, Weinstock GM, Davidson EH *et al.*, 2006. The genome of the sea urchin *Strongylocentrotus purpuratus*. Science 314: 941-952.

Sogin EM, Putnam HM, Anderson PE, Gates RD. 2016. Metabolomic signatures of increases in temperature and ocean acidification from the reef-building coral, *Pocillopora damicornis*. Metabolomics 12:71.

Somero GN. 2010. The physiology of climate change: how potentials for acclimatization and genetic adaptation will determine "winners" and "losers." J Exp Biol 213:912–20.

Strathmann MF. 2017. Reproduction and Development of Marine Invertebrates of the Northern Pacific Coast: Data and Methods for the Study of Eggs, Embryos, and Larvae University of Washington Press.

Stumpp M, Hu MY, Melzner F, Gutowska MA, Dorey N, Himmerkus N, Holtmann WC, Dupont ST, Thorndyke MC, Bleich M. 2012. Acidified seawater impacts sea urchin larvae pH regulatory systems relevant for calcification. Proc Natl Acad Sci U S A 109:18192–97.

Sunday JM, Calosi P, Dupont S, Munday PL, Stillman JH, Reusch TBH. 2014. Evolution in an acidifying ocean. Trends Ecol Evol 29:117–25.

Suwa R, Nojiri Y, Ono T, Shirayama Y. 2013. Effects of low pCO₂ conditions on sea urchin larval size. Mar Ecol 34:443–50.

Vázquez DP, Gianoli E, Morris WF, Bozinovic F. 2017. Ecological and evolutionary impacts of changing climatic variability. Biol Rev Camb Philos Soc 92:22–42.

Vidavsky N, Addadi S, Mahamid J, Shimoni E, Ben-Ezra D, Shpigel M, Weiner S, Addadi L. 2014. Initial stages of calcium uptake and mineral deposition in sea urchin embryos. Proc Natl Acad Sci U S A 111:39–44.

Vlachos C, Burny C, Pelizzola M, Borges R, Futschik A, Kofler R, Schlötterer C. 2019.

Benchmarking software tools for detecting and quantifying selection in evolve and resequencing studies. Genome Biol 20:169.

Yu G, Li F, Qin Y, Bo X, Wu Y, Wang S. 2010. GOSemSim: an R package for measuring semantic similarity among GO terms and gene products. Bioinformatics 26:976–78.

Yu PC, Matson PG, Martz TR, Hofmann GE. 2011. The ocean acidification seascape and its relationship to the performance of calcifying marine invertebrates: Laboratory experiments on the development of urchin larvae framed by environmentally-relevant pCO₂/pH. Journal of Experimental Marine Biology and Ecology.